arXiv:1107.0295v1 [math.AG] 1 Jul 2011

Direct calculation of higher rank invariants of stable pairs using the Joyce machinery of stack functions

Artan Sheshmani

July 4, 2011

Abstract

The rank 2 invariants of stable pairs over a Calabi-Yau threefold were computed in [15] using the wall-crossing formula of Joyce-Song [9] and Kontsevich-Soibelman [12]. The computations in [15] depended on the combinatorial properties of the elements of the Hall-algebra given as stack functions defined by Joyce [4]. These combinatorial computations become immediately complicated and hard to carry out when studying higher rank stable pairs with rank>2. In this article we introduce an independent approach to computation of rank 2 stable pair invariants not using the wallcrossing formula and rather using the stratification of their corresponding moduli spaces and directly computing the weighted Euler characteristic of the strata. This approach may be used to avoid complex combinatorial wallcrossing calculations and to carry out the computation of invariants in rank>2 cases directly. Moreover, we suspect that the method introduced in this article may be used to prove the integrality of the corresponding partition functions for higher rank invariants in special cases.

Contents

1 Introduction **2**

2 Acknowledgment **3**

3 Preliminary Definitions **4**
3.1 Definition of frozen and highly frozen triples 4
3.2 The auxiliary category $\mathcal{B}_p$ 6

4 Moduli stack of objects in $\mathcal{B}_p$ **8**
4.1 The moduli stacks $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_{p,ss}}(\tilde{\tau})$ and $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}^{\mathbf{R}}_{p,ss}}(\tilde{\tau})$ 10

5 Stack function identities in Ringel Hall algebra **11**

6 Direct computation of invariants in an example **13**
6.1 Stacky structure of $\mathfrak{M}^{([\mathbb{P}^1],2)}_{sp,\mathcal{B}_p}(\tilde{\tau})$ 14
6.2 Stacky structure of $\mathfrak{M}^{([\mathbb{P}^1],2)}_{nsp,\mathcal{B}_p}(\tilde{\tau})$ 15
6.3 Computation of $\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau})*\bar{\delta}_s^{(0,1)}(\tilde{\tau})$ 16

6.4 Computation of $\bar{\delta}_s^{(0,1)}(\tilde{\tau}) * \bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau})$ . . . 18
6.5 Computation of $\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})$ . . . 18
6.6 Computation of the invariant . . . 19

7 An approach to direct calculations in general cases **20**

8 $\mathrm{GL}_2(\mathbb{C})$**-invariant stratification** **22**

9 Computation of $\bar{\delta}_s^{(\beta_k,1)}(\tilde{\tau}) * \bar{\delta}_s^{(\beta_l,1)}(\tilde{\tau})$ **31**
9.1 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_1^{(k,l)}(\tilde{\tau}))$ . . . 35
9.2 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))$ . . . 37
9.3 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))$ . . . 41
9.4 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(k,l)}(\tilde{\tau}))$ . . . 45
9.5 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))$ . . . 45
9.6 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{(\beta,2)}(\tilde{\tau}))$ . . . 46

10 Computation of the invariant **47**

1 Introduction

In [14] we introduced the highly frozen triples and computed their invariants using the Graber-Pandharipande [2] method of virtual localization over the local Calabi-Yau threefolds such as the total space of $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1) \to \mathbb{P}^1$. The objects studied in [14] (because of the stability condition chosen) were reminiscent of the higher rank analogue of the Pandharipande-Thomas stable pairs [13] (Section 12.1). In [15] we used the machinery of Joyce-Song [9] and Kontsevich-Soibelman [12] in wallcrossings to compute the invariants of the objects of an auxiliary category $\mathcal{B}_p$ [15] (Definition 4.2 and Section 5) in terms of the generalized Donaldson-Thomas invariants. Moreover, we proved [15] (Section 13) that these auxiliary objects are related to highly frozen triples equipped with a new stability condition which turned them in to an analogue of the higher rank Joyce-Song stable pairs [9]. As a result we found that the invariants of the rank 2 semistable Joyce-Song pairs are related to the generalized Donaldson-Thomas invariants by the following identity [15] (Equation 12.32):

$$\begin{aligned}&\mathrm{HFT}(X, P_\beta, r, \hat{\tau}) = \chi(\mathfrak{M}_{s,\mathrm{HFT}}^{(P_\beta,r)}(\hat{\tau}), \nu_{\mathfrak{M}_{\mathrm{HFT}}}^{(P_\beta,r)}) = \chi(\mathfrak{M}_{\mathcal{B}_p,ss}^{(\beta,r)}(\tilde{\tau}), \nu_{\mathfrak{M}_{\mathcal{B}_p,ss}}^{(\beta,r)}) = \\ &\sum_{1\leq l,\beta_1+\cdots+\beta_l=\beta} \frac{-1}{4} \cdot \left[\frac{(1)}{l!} \cdot \prod_{i=1}^{l} \left(\overline{DT}^{\beta_i}(\tau) \cdot \bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots+\beta_{i-1},2),(\beta_i,0))\right.\right. \\ &\left.\left. \cdot (-1)^{\bar{\chi}_{\mathcal{B}_p}((0,2),(\beta_1,0))+\sum_{i=1}^{l}\bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots\beta_{i-1},2),(\beta_i,0))}\right)\right]. \qquad (1.1)\end{aligned}$$

According to [15] (Corollary 13.7) and via carrying out computational examples we conjecture that for the case of rank> 2 the invariants of the higher rank Joyce-Song stable pairs can be computed in terms of the generalized Donaldson-Thomas invariants by the following identity:

$$\begin{aligned}&\mathrm{HFT}(X,P_\beta,r,\hat{\tau})=\chi(\mathfrak{M}^{(P_\beta,r)}_{s,\mathrm{HFT}}(\hat{\tau}),\nu^{(P_\beta,r)}_{\mathfrak{M}_{\mathrm{HFT}}})=(-1)^{r^2}\cdot\chi(\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_p,ss}(\tilde{\tau}),\nu^{(\beta,r)}_{\mathfrak{M}_{\mathcal{B}_{p,ss}}})=\\&(-1)^{r^2+1}\cdot\sum_{1\leq l,\beta_1+\cdots+\beta_l=\beta}\frac{1}{r^2}\cdot\left[\frac{(1)}{l!}\cdot\prod_{i=1}^{l}\left(\overline{DT}^{\beta_i}(\tau)\cdot\bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots+\beta_{i-1},r),(\beta_i,0))\right.\right.\\&\left.\left.\cdot(-1)^{\bar{\chi}_{\mathcal{B}_p}((0,r),(\beta_1,0))+\sum_{i=1}^{l}\bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots\beta_{i-1},r),(\beta_i,0))}\right)\right],\end{aligned}\tag{1.2}$$

where r denotes the rank of the highly frozen triples. The purpose of this article is to introduce a *direct* method of calculation for the invariants of the higher rank Joyce-Song pairs in terms of the generalized Donaldson-Thomas invariants for rank= 2 cases. Here by direct we mean a calculation not involving the wallcrossing computation as in [15] and rather involving the stratification of the moduli stack of the auxiliary objects in the category $\mathcal{B}_p$, computing the weighted Euler characteristics of the strata and finally exploiting [15] (Corollary 13.7) to obtain an identity which relates the obtained weighted Euler characteristics of the strata to the invariants of the higher rank Joyce-Song stable pairs and the generalized Donaldson-Thomas invariants. We show in this article that the result of our calculation agrees with that of [15] (Equation 12.32).

We believe that following the same computational strategy would enable one to compute the invariants for the case of rank> 2 and to obtain Equation (1.2) while avoiding the complicated combinatorial calculations involved in the wallcrossing computations. Moreover, we suspect that the method introduced in this article may be used to prove the integrality conjectures for the partition functions of the higher rank Joyce-Song invariants in special cases. Toda in [16] has used a similar stratification technique and has provided an evidence of the integrality conjecture proposed by Kontsevich-Soibelman [12] (Conjecture 6).

2 Acknowledgment

I thank my adviser Sheldon Katz for his constant help and guidance over the past years and through this project and for suggesting me the computation in Section 6. I would like to thank Yukinobu Toda for many valuable discussions and for explaining me his paper on rank= 2 Donaldson-Thomas invariants. I thank Arend Bayer for kindly answering some of my questions related to this work in spring 2011. I am grateful to Richard Thomas for his help and support and for providing me the opportunity for being a member at Isaac Newton Institute for Mathematical sciences during spring 2011. I also thank Newton Institute for their kind hospitality during the time that I was visiting. The stimulating environment at Newton Institute and many productive conversations with experts helped me to fix some of the weaknesses which existed in the early stages of this article. I acknowledge support from NSF grants DMS 0244412, DMS 0555678 and DMS 08-38434 EMSW21-MCTP (R.E.G.S).

3 Preliminary Definitions

3.1 Definition of frozen and highly frozen triples

In this section we borrow some of the definitions in [15] (Section 3) which are crucial for our computations in the later sections.

Definition 3.1. Let X be a nonsingular projective Calabi-Yau 3-fold over $\mathbb{C}$ (i.e $K_X \cong \mathcal{O}_X$ and $\pi_1(X) = 0$ which implies $\mathrm{H}^1(\mathcal{O}_X) = 0$) with a fixed polarization L . A holomorphic triple supported over X is given by (E_1, E_2, ϕ) consisting of a torsion free coherent sheaf E_1 and a pure sheaf with one dimensional support E_2, together with a holomorphic morphism $\phi : E_1 \to E_2$.

A homomorphism of triples from $(\acute{E}_1, \acute{E}_2, \acute{\phi})$ to (E_1, E_2, ϕ) is a commutative diagram:

$$\begin{tikzcd}
\acute{E}_1 \arrow[r, "\acute{\phi}"] \arrow[d] & \acute{E}_2 \arrow[d] \\
E_1 \arrow[r, "\phi"] & E_2
\end{tikzcd}$$

Remark 3.2. A triple (E_1, E_2, ϕ) of type (P_1, P_2, β) is given by a triple such that $P(E_1(m)) = P_1$ and $P(E_2(m)) = P_2$ and $\beta = \mathrm{ch}_2(E_2)$ as defined in Definition 3.8. During the discussion, for simplicity, we omit β and write a triple of type (P_1, P_2).

Remark 3.3. Since by assumption the sheaf E_2 has one dimensional support, the Hilbert polynomial of E_2 in variable m satisfies:

$$P(E_2(m)) = \chi(E_2(m)) = m \int_{\beta} c_1(L) + d. \tag{3.1}$$

Here $c_1(L)$ is the first Chern class of the fixed polarization L over X and $d \in \mathbb{Z}$ and β as before is $\mathrm{ch}_2(E_2)$. Note that P_2 is a polynomial of degree= $\dim(\mathrm{Supp}(E_2)) = 1$ and by rank of E_2 (denoted by $rk(E_2)$) we mean the leading coefficient of P_2.

Definition 3.4. A *frozen triple* of rank r is a special case of a holomorphic triple where $E_1 \cong \mathcal{O}_X(-n)^{\oplus r}$ for some $n \in \mathbb{Z}$.

Remark 3.5. By freezing the triple we mean fixing E_1 to be isomorphic with $\mathcal{O}_X(-n)^{\oplus r}$. We do not make a choice of such an isomorphism here. Later we fix an isomorphism $\psi : E_1 \xrightarrow{\cong} \mathcal{O}_X(-n)^{\oplus r}$and we call the triples *highly-frozen* triples.

Definition 3.6. Use the notation above. Let S be a $\mathbb{C}$ scheme of finite type and let $\pi_X : X \times S \to X$ and $\pi_S : X \times S \to S$ be the corresponding projections. An *S-flat family* of triples over X is a triple $(\mathcal{E}_1, \mathcal{E}_2, \phi)$ consisting of a morphism of $\mathcal{O}_{X\times S}$ modules $\mathcal{E}_1 \xrightarrow{\phi} \mathcal{E}_2$ such that $\mathcal{E}_1$ and $\mathcal{E}_2$ are flat over S and for every point $s \in S$ the fiber $(\mathcal{E}_1, \mathcal{E}_2, \phi) \mid_s$ is given by a holomorphic triple as in Definitions 3.1.

Two S-flat families of triples $(\mathcal{E}_1, \mathcal{E}_2, \phi)$ and $(\mathcal{E}'_1, \mathcal{E}'_2, \phi')$ are isomorphic if there exists a commutative diagram of the form:

$$\begin{tikzcd}
\acute{\mathcal{E}}_1 \arrow[r, "\acute{\phi}"] \arrow[d, "\cong"'] & \acute{\mathcal{E}}_2 \arrow[d, "\cong"] \\
\mathcal{E}_1 \arrow[r, "\phi"] & \mathcal{E}_2
\end{tikzcd}$$

Definition 3.7. An S-flat family of frozen-triples is a triple $(\mathcal{E}_1, \mathcal{E}_2, \phi)$ consisting of a morphism of $\mathcal{O}_{X\times S}$ modules $\phi : \mathcal{E}_1 \to \mathcal{E}_2$ such that $\mathcal{E}_1$ and $\mathcal{E}_2$ satisfy the condition of Definition 3.6 and moreover $\mathcal{E}_1 \cong \pi_X^*\mathcal{O}_X(-n) \otimes \pi_S^*\mathcal{M}_S$ where $\mathcal{M}_S$ is a vector bundle of rank r on S.
Two S-flat families of frozen-triples $(\mathcal{E}_1, \mathcal{E}_2, \phi)$ and $(\mathcal{E}'_1, \mathcal{E}'_2, \phi')$ are isomorphic if there exists a commutative diagram:

$$\begin{tikzcd}
\acute{\mathcal{E}}_1 \arrow[r, "\acute{\phi}"] \arrow[d, "\cong"'] & \acute{\mathcal{E}}_2 \arrow[d, "\cong"] \\
\mathcal{E}_1 \arrow[r, "\phi"] & \mathcal{E}_2
\end{tikzcd}$$

Definition 3.8. A frozen-triple of class β and of fixed Hilbert polynomial P_2 is a frozen-triple (E_1, E_2, ϕ) such that the Hilbert polynomial of E_2 is equal to P_2 and $\beta = \mathrm{ch}_2(E_2)$. Having fixed r in $E_1 \cong \mathcal{O}_X^{\oplus r}(-n)$, we denote these frozen triples as frozen triples of type (P_2, r).

Now we define highly frozen triples.
Definition 3.9. A *highly frozen triple* is a quadruple (E_1, E_2, ϕ, ψ) where (E_1, E_2, ϕ) is a frozen triple as in Definition 3.4 and $\psi : E_1 \xrightarrow{\cong} \mathcal{O}_X(-n)^{\oplus r}$ is a fixed choice of isomorphism. A morphism between highly frozen triples $(E'_1, E'_2, \phi', \psi')$ and (E_1, E_2, ϕ, ψ) is a morphism $E'_2 \xrightarrow{\rho} E_2$ such that the following diagram is commutative.

$$\begin{tikzcd}
\mathcal{O}_X(-n)^{\oplus r} \arrow[r, "\psi'^{-1}"] \arrow[d, "id"'] & E'_1 \arrow[r, "\acute{\phi}"] \arrow[d] & E'_2 \arrow[d, "\rho"] \\
\mathcal{O}_X(-n)^{\oplus r} \arrow[r, "\psi^{-1}"] & E_1 \arrow[r, "\phi"] & E_2
\end{tikzcd}$$

Definition 3.10. An S-flat family of highly frozen-triples is a quadruple $(\mathcal{E}_1, \mathcal{E}_2, \phi, \psi)$ consisting of a morphism of $\mathcal{O}_{X\times S}$ modules $\mathcal{E}_1 \xrightarrow{\phi} \mathcal{E}_2$ such that $\mathcal{E}_1$ and $\mathcal{E}_2$ satisfy the condition of Definition 3.6 and moreover $\psi : \mathcal{E}_1 \xrightarrow{\cong} \pi_X^*\mathcal{O}_X(-n) \otimes \pi_S^*\mathcal{O}_S^{\oplus r}$ is a fixed choice of isomorphism.
Two S-flat families of highly frozen-triples $(\mathcal{E}_1, \mathcal{E}_2, \phi, \psi)$ and $(\mathcal{E}'_1, \mathcal{E}'_2, \phi', \psi')$ are isomorphic if there exists a commutative diagram:

$$\begin{tikzcd}
\pi_X^*\mathcal{O}_X(-n) \otimes \pi_S^*\mathcal{O}_S^{\oplus r} \arrow[r, "\psi'^{-1}"] \arrow[d, "id"'] & \mathcal{E}'_1 \arrow[r, "\acute{\phi}"] \arrow[d] & \mathcal{E}'_2 \arrow[d, "\cong"] \\
\pi_X^*\mathcal{O}_X(-n) \otimes \pi_S^*\mathcal{O}_S^{\oplus r} \arrow[r, "\psi^{-1}"] & \mathcal{E}_1 \arrow[r, "\phi"] & \mathcal{E}_2
\end{tikzcd}$$

Definition 3.11. Use notation in Definitions 3.1 to 3.7. Given a coherent sheaf E_2, let p_{E_2} denote the reduced Hilbert polynomial of E_2 with respect to the ample line bundle $\mathcal{O}_X(1)$. A highly frozen

triple, (E_1, E_2, ϕ, ψ), given by the data $E_1 \xrightarrow{\phi} E_2$ and $\psi : \mathcal{O}_X(-n) \xrightarrow{\cong} E_1$, is called $q \to 0$ $\hat{\tau}$-limit stable (or in short $\hat{\tau}$-limit stable) if:
1. $p_{\acute{E}_2} \leq p_{E_2}$ for all proper subsheaves $\acute{E}_2$ such that $\acute{E}_2 \neq 0$.
2. If ϕ factors through $\acute{E}_2$ ($\acute{E}_2$ a proper subsheaf of E_2), then $p_{\acute{E}_2} < p_{E_2}$.
Here the same notion of stability can be applied to frozen triples.

3.2 The auxiliary category $\mathcal{B}_p$

In this section we give the definition and the properties of the objects in the auxiliary category $\mathcal{B}_p$. As was shown in [15] (Section 13), the objects in $\mathcal{B}_p$ (when equipped with the right notion of semistability) are closely related to $\hat{\tau}$-limit stable highly frozen triples.

Definition 3.12. (Joyce and Song) [9] (Definition 13.1). Let X be a Calabi-Yau threefold equipped with ample line bundle $\mathcal{O}_X(1)$. Let τ denote the Giseker stability on the abelian category of coherent sheaves supported over X. Define $\mathcal{A}_p$ to be the sub-category of coherent sheaves whose objects are zero sheaves and non-zero τ-semistable sheaves with reduced Hilbert polynomial p.

Definition 3.13. (Joyce and Song) [9]. Define category $\mathcal{B}_p$ to be the category whose objects are triples (F, V, ϕ), where $F \in Obj(\mathcal{A}_p)$, V is a finite dimensional $\mathbb{C}$-vector space, and $\phi : V \to \operatorname{Hom}(\mathcal{O}_X(-n), F)$ is a $\mathbb{C}$-linear map. Given (F, V, ϕ) and $(\acute{F}, \acute{V}, \acute{\phi})$ in $\mathcal{B}_p$ define morphisms $(F, V, \phi) \to (\acute{F}, \acute{V}, \acute{\phi})$ in $\mathcal{B}_p$ to be pairs of morphisms (f, g) where $f : F \to \acute{F}$ is a morphism in $\mathcal{A}_p$ and $g : V \to \acute{V}$ is a $\mathbb{C}$-linear map, such that the following diagram commutes:

$$\begin{tikzcd}
V \arrow[r, "\phi"] \arrow[d, "g"'] & \operatorname{Hom}(\mathcal{O}_X(-n), F) \arrow[d, "f"] \\
\acute{V} \arrow[r, "\acute{\phi}"] & \operatorname{Hom}(\mathcal{O}_X(-n), \acute{F})
\end{tikzcd} \tag{3.2}$$

Now we define the numerical class of objects in $\mathcal{B}_p$:

Definition 3.14. Let $\mathcal{A}$ denote any abelian category. Let $\mathcal{K}_0(\mathcal{A})$ denote the Grothendieck group of $\mathcal{A}$ generated by isomorphism classes $[E]$ of objects E in $\mathcal{A}$ which satisfy the relation $[E] = [F] + [G]$ if there exists a short exact sequence $0 \to F \to E \to G \to 0$ in $\mathcal{A}$. The Euler form $\bar{\chi} : \mathcal{K}_0(\mathcal{A}) \times \mathcal{K}_0(\mathcal{A}) \to \mathbb{Z}$ is defined as:

$$\bar{\chi}([E], [F]) = \sum_{i \geq 0} (-1)^i \dim Ext^i(E, F).$$

Definition 3.15. Let $I = \{\alpha \in \mathcal{K}_0(\mathcal{A}) \mid \bar{\chi}(\alpha, \beta) = \bar{\chi}(\beta, \alpha) = 0, \forall \beta \in \mathcal{K}_0(\mathcal{A})\}$. Define the numerical Grothendieck group of $\mathcal{A}$ to be the quotient of $\mathcal{K}_0(\mathcal{A})$ by the two sided kernel of $\bar{\chi}$, i.e $\mathcal{K}^{num}(\mathcal{A}) = \mathcal{K}_0(\mathcal{A})/I$. Define the positive cone of $\mathcal{A}$, $C(\mathcal{A}) \subset \mathcal{K}(\mathcal{A}) = \mathcal{K}^{num}(\mathcal{A})$ to be

$$C(\mathcal{A}) = \{[E] \in K(\mathcal{A}) \mid E \ncong 0, E \in \mathcal{A}\}. \tag{3.3}$$

The above definitions extend to the case where the abelian category is $\mathcal{A}_p$:

Definition 3.16. Define $\mathcal{K}(\mathcal{B}_p) = \mathcal{K}(\mathcal{A}_p) \oplus \mathbb{Z}$ where for $(F, V, \phi) \in \mathcal{B}_p$, $[(F, V, \phi)] = ([F], \dim(V))$.

We state following results by Joyce and Song without proof.

Lemma 3.17. *(Joyce and Song) [9] (Lemma 13.2). The category $\mathcal{B}_p$ is abelian and $\mathcal{B}_p$ satisfies the condition that for the underlying sheaves F the following is true:*
If $[F] = 0 \in \mathcal{K}^{num}(\mathcal{A}_p)$ then $F \cong 0$, moreover $\mathcal{B}_p$ is noetherian and artinian and the moduli stacks $\mathfrak{M}_{\mathcal{B}_p}^{(\beta,d)}$ are of finite type $\forall(\beta, d) \in C(\mathcal{B}_p)$.

Remark 3.18. The category $\mathcal{A}_p$ embeds as a full and faithful sub-category in $\mathcal{B}_p$ by $F \to (F, 0, 0)$, moreover it is shown by Joyce and Song in [9] that every object (F, V, ϕ) sits in a short exact sequence.

$$0 \to (F, 0, 0) \to (F, V, \phi) \to (0, V, 0) \to 0 \tag{3.4}$$

Definition 3.19. (Joyce and Song) [9]. Define $\bar{\chi}^{\mathcal{B}_p} : \mathcal{K}^{num}(\mathcal{B}_p) \times \mathcal{K}^{num}(\mathcal{B}_p) \to \mathbb{Z}$ by:

$$\begin{aligned} \bar{\chi}^{\mathcal{B}_p}((\beta, d), (\gamma, e)) &= \bar{\chi}(\beta - d[\mathcal{O}_X(-n)], \gamma - e[\mathcal{O}_X(-n)]) \\ &= \bar{\chi}(\beta, \gamma) - d\bar{\chi}([\mathcal{O}_X(-n)], \gamma) + e\bar{\chi}([\mathcal{O}_X(-n)], \beta) \end{aligned} \tag{3.5}$$

Definition 3.20. (Joyce and Song) [9] (Definition. 13.5). Define the positive cone of $\mathcal{B}_p$ by:

$$\mathcal{C}(\mathcal{B}_p) = \{(\beta, d) \mid \beta \in \mathcal{C}(\mathcal{A}_p) \text{ and } d \geq 0 \text{ or } \beta = 0 \text{ and } d > 0\}.$$

Next we recall the definition of weak (semi)stability from [9] for a general abelian category $\mathcal{A}$.

Definition 3.21. (Joyce and Song)[9](Definition. 3.5). Let $\mathcal{A}$ be an abelian category, $\mathcal{K}(\mathcal{A})$ be the quotient of $\mathcal{K}_0(\mathcal{A})$ by some fixed subgroup, and $C(\mathcal{A})$ the positive cone of $\mathcal{A}$. Suppose $(T, \leq)$ is a totally ordered set and $\tau : C(\mathcal{A}) \to T$ a map. We call $(\tau, T, \leq)$ a stability condition on $\mathcal{A}$ if whenever $\alpha, \beta, \gamma \in C(\mathcal{A})$ with $\beta = \alpha + \gamma$ then either $\tau(\alpha) < \tau(\beta) < \tau(\gamma)$ or $\tau(\alpha) > \tau(\beta) > \tau(\gamma)$ or $\tau(\alpha) = \tau(\beta) = \tau(\gamma)$. We call $(\tau, T, \leq)$ a weak stability condition on $\mathcal{A}$ if if whenever $\alpha, \beta, \gamma \in C(\mathcal{A})$ with $\beta = \alpha + \gamma$ then either $\tau(\alpha) \leq \tau(\beta) \leq \tau(\gamma)$ or $\tau(\alpha) \geq \tau(\beta) \geq \tau(\gamma)$.
For such $(\tau, T, \leq)$, we say that a nonzero object E in $\mathcal{A}$ is

1. τ-semistable if $\forall S \subset E$ where $S \ncong 0$, we have $\tau([S]) \leq \tau([E/S])$
2. τ-stable if $\forall S \subset E$ where $S \ncong 0$, we have $\tau([S]) < \tau([E/S])$
3. τ-unstable if it is not semistable.

In our analysis we apply the Definition 3.21 to objects in $\mathcal{A}_p$. Next we define the notion of permissible stability condition from [9] for the moduli stack of objects in a general abelian category $\mathcal{A}$. Here we assume that the moduli stack of τ-(semi)stable objects in $\mathcal{A}$ exists.

Definition 3.22. (Joyce and Song) [9] (Definition. 3.7). Let $(\tau, T, \leq)$ be a weak stability condition on $\mathcal{A}$. For $\alpha \in \mathcal{K}(\mathcal{A})$ let $\mathfrak{M}_{ss}^{\alpha}$ and $\mathfrak{M}_{st}^{\alpha}$ denote the moduli stacks of τ-(semi)stable objects $E \in \mathcal{A}$ with class $[E] = \alpha$ in $\mathcal{K}(\mathcal{A})$. We call $(\tau, T, \leq)$ permissible if

1. $\mathcal{A}$ is τ-Artinian (There exists no infinite chains of sub-objects with reducing slope of the subsequent quotients).
2. $\mathfrak{M}_{ss}^{\alpha}(\tau)$ is a finite type sub-stack of $\mathfrak{M}_{\mathcal{A}}^{\alpha}$ $\forall \alpha \in C(\mathcal{A})$.

One example of an abelian category for which there exists a moduli stack of (semi)stable objects which satisfy the condition in Definition 3.22 is the category of coherent sheaves, $\mathrm{Coh}(X)$. In that case τ is the Giseker stability condition and from usual arguments, it is clear that there exist finite type open substacks $\mathfrak{M}_{s}^{\alpha}(\tau)$ and $\mathfrak{M}_{ss}^{\alpha}(\tau)$ of τ stable and τ-semistable sheaves $F \in \mathcal{A}$ with numerical class $\alpha \in \mathcal{K}^{num}(\mathrm{Coh}(X))$.

Definition 3.23. (Joyce and Song) [9] (Definition. 13.5). Define the weak stability condition $\tilde{\tau}$ in $\mathcal{B}_p$ by:
1. $\tilde{\tau}(\beta, d) = 0$ if $d = 0$ and $\tilde{\tau}(\beta, d) = 1$ if $d > 0$.

4 Moduli stack of objects in $\mathcal{B}_p$

In this section for the purpose of completeness, we review the definition and construction of the moduli stack of objects in $\mathcal{B}_p$ from [15] (sections 5, 5.1 and 5.2).

Remark 4.1. By [9] (Page 185) there exists a natural embedding functor $\mathfrak{F} : \mathcal{B}_p \to D(X)$ which takes $(F, V, \phi_V) \in \mathcal{B}_p$ to an object in the derived category given by $\cdots \to 0 \to V \otimes \mathcal{O}_X(-n) \to F \to 0 \to \cdots$ where $V \otimes \mathcal{O}_X(-n)$ and F sit in degree -1 and 0. Assume that $\dim(V) = r$. In that case $V \otimes \mathcal{O}_X(-n) \cong \mathcal{O}_X(-n)^{\oplus r}$. Hence one may view an object $(F, V, \phi_V) \in \mathcal{B}_p$ as a triple (E, F, ϕ) represented by a complex $\phi : E \to F$ such that $E \cong \mathcal{O}_X(-n)^{\oplus r}$ (note the similarity between the objects in $\mathcal{B}_p$ and frozen triples in Definition 3.4).
Definition 4.2. Fix a parametrizing scheme of finite type S. Let $\pi_X : X \times S \to X$ and $\pi_S : X \times S \to S$ denote the natural projections. Use the natural embedding functor $\mathfrak{F} : \mathcal{B}_p \to D(X)$ [9] (Page 185). Define the S-flat family of objects in $\mathcal{B}_p$ of type (β, r) as a complex

$$\pi_S^* M \otimes \pi_X^* \mathcal{O}_X(-n) \xrightarrow{\psi_S} \mathcal{F}$$

sitting in degree -1 and 0 such that $\mathcal{F}$ is given by an S-flat family of semistable sheaves with fixed reduced Hilbert polynomial p with $\mathrm{Ch}(F) = \beta$ and M is a vector bundle of rank r over S. A morphism between two such S-flat families is given by a morphism between the complexes $\pi_S^* M \otimes \pi_X^* \mathcal{O}_X(-n) \xrightarrow{\psi_S} \mathcal{F}$ and $\pi_S^* M' \otimes \pi_X^* \mathcal{O}_X(-n) \xrightarrow{\psi'_S} \mathcal{F}'$:

$$\begin{tikzcd}
\pi_S^* M \otimes \pi_X^* \mathcal{O}_X(-n) \arrow[r, "\psi_S"] \arrow[d] & \mathcal{F} \arrow[d] \\
\pi_S^* M' \otimes \pi_X^* \mathcal{O}_X(-n) \arrow[r, "\psi'_S"] & \mathcal{F}'.
\end{tikzcd}$$

Moreover an isomorphism between two such S-flat families in $\mathcal{B}_p$ is given by an isomorphism between the associated complexes $\pi_S^* M \otimes \pi_X^* \mathcal{O}_X(-n) \xrightarrow{\psi_S} \mathcal{F}$ and $\pi_S^* M' \otimes \pi_X^* \mathcal{O}_X(-n) \xrightarrow{\psi'_S} \mathcal{F}'$:

$$\begin{tikzcd}
\pi_S^* M \otimes \pi_X^* \mathcal{O}_X(-n) \arrow[r, "\psi_S"] \arrow[d, "\cong"'] & \mathcal{F} \arrow[d, "\cong"] \\
\pi_S^* M' \otimes \pi_X^* \mathcal{O}_X(-n) \arrow[r, "\psi'_S"] & \mathcal{F}'.
\end{tikzcd}$$

Note the similarity between definition of isomorphism between S-flat families of objects of type (β, r) in $\mathcal{B}_p$ and the isomorphism between two S-flat families of frozen triples of type (P_2, r) in

Definition 3.7.

From now on whenever we mention objects in $\mathcal{B}_p$ we mean the objects which lie in the image of the natural embedding functor $\mathfrak{F} : \mathcal{B}_p \to D(X)$ [9] (Page 185). Moreover by the S-flat family of objects in $\mathcal{B}_p$, their morphisms (or isomorphisms) we mean the corresponding definitions as in Definition 4.2. Now we define the *rigidified* objects in $\mathcal{B}_p$. We give the category of these objects a new name $\mathcal{B}_p^{\mathbf{R}}$. However, we emphasize that it is implicitly understood that for us the category $\mathcal{B}_p^{\mathbf{R}}$ is the same as $\mathcal{B}_p$ together with an additional structure:

Definition 4.3. Define the category $\mathcal{B}_p^{\mathbf{R}}$ to be the category of rigidified objects in $\mathcal{B}_p$ whose objects are defined by tuples $(F, \mathbb{C}^{\oplus r}, \rho)$ where F is a coherent sheaf with reduced Hilbert polynomial p and $\mathrm{Ch}(F) = \beta$ and $\rho : \mathbb{C}^r \to \mathrm{Hom}(\mathcal{O}_X(-n), F)$ (for some r). Given two rigidified objects of fixed given type (β, r) as $(F, \mathbb{C}^{\oplus r}, \rho)$ and $(F', \mathbb{C}^{\oplus r}, \rho')$ in $\mathcal{B}_p^{\mathbf{R}}$ define morphisms $(F, \mathbb{C}^{\oplus r}, \rho) \to (F', \mathbb{C}^{\oplus r}, \rho')$ to be given by a morphism $f : F \to F'$ in $\mathcal{A}_p$ such that the following diagram commutes:

$$\begin{tikzcd}
\mathbb{C}^{\oplus r} \arrow[r, "\rho"] \arrow[d, "\mathrm{id}"'] & \mathrm{Hom}(\mathcal{O}_X(-n), F) \arrow[d, "f"] \\
\mathbb{C}^{\oplus r} \arrow[r, "\rho'"] & \mathrm{Hom}(\mathcal{O}_X(-n), \acute{F}).
\end{tikzcd} \tag{4.1}$$

Remark 4.4. There exists a natural embedding functor $\mathfrak{F}^{\mathbf{R}} : \mathcal{B}_p^{\mathbf{R}} \to D(X)$ which takes $(F, \mathbb{C}^{\oplus r}, \rho) \in \mathcal{B}_p^{\mathbf{R}}$ to an object in the derived category given by $\cdots \to 0 \to \mathbb{C}^{\oplus r} \otimes \mathcal{O}_X(-n) \to F \to 0 \to \cdots$ where $\mathbb{C}^{\oplus r} \otimes \mathcal{O}_X(-n)$ sits in degree -1 and F sits in degree 0. One may view an object in $\mathcal{B}_p^{\mathbf{R}}$ as a quadruple (E, F, ϕ, ψ) represented by a complex $\phi : E \to F$ such that $\psi : E \cong \mathcal{O}_X(-n)^{\oplus r}$ is a fixed choice of isomorphism (note the similarity between the objects in $\mathcal{B}_p^{\mathbf{R}}$ and highly frozen triples in Definition 3.9).

Definition 4.5. Fix a parametrizing scheme of finite type S. Let $\pi_X : X \times S \to X$ and $\pi_S : X \times S \to S$ denote the natural projections. Use the natural embedding functor $\mathfrak{F}^{\mathbf{R}} : \mathcal{B}_p^{\mathbf{R}} \to D(X)$ in Remark 4.4. An S-flat family of objects of type (β, r) in $\mathcal{B}_p^{\mathbf{R}}$ is given by a complex

$$\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \xrightarrow{\psi_S} \mathcal{F}$$

sitting in degree -1 and 0 such that $\mathcal{F}$ is given by an S-flat family of semistable sheaves with fixed reduced Hilbert polynomial p with $\mathrm{Ch}(F) = \beta$. A morphism between two such S-flat families in $\mathcal{B}_p^{\mathbf{R}}$ is given by a morphism between the complexes $\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \xrightarrow{\psi_S} \mathcal{F}$ and $\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \xrightarrow{\psi'_S} \mathcal{F}'$:

$$\begin{tikzcd}
\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \arrow[r, "\psi_S"] \arrow[d, "\mathrm{id}_{\mathcal{O}_{X\times S}}"'] & \mathcal{F} \arrow[d] \\
\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \arrow[r, "\psi'_S"] & \mathcal{F}'.
\end{tikzcd}$$

Moreover an isomorphism between two such S-flat families in $\mathcal{B}_p^{\mathbf{R}}$ is given by an isomorphism between the associated complexes $\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \xrightarrow{\psi_S} \mathcal{F}$ and $\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \xrightarrow{\psi'_S} \mathcal{F}'$:

$$\begin{tikzcd}
\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \arrow[r, "\psi_S"] \arrow[d, "\mathrm{id}_{\mathcal{O}_{X\times S}}"'] & \mathcal{F} \arrow[d, "\cong"] \\
\pi_S^*\mathcal{O}_S^{\oplus r} \otimes \pi_X^*\mathcal{O}_X(-n) \arrow[r, "\psi'_S"] & \mathcal{F}'.
\end{tikzcd}$$

Note the similarity between definition of isomorphism between S-flat families of objects of type (β, r) in $\mathcal{B}_p^{\mathbf{R}}$ and the isomorphism between two S-flat families of highly frozen triples of type (P_2, r) in definitions 3.9 and 3.10.

Similar to the way that we treated objects in $\mathcal{B}_p$ from now on whenever we mention objects in $\mathcal{B}_p^{\mathbf{R}}$ we mean the objects which lie in the image of the natural embedding functor $\mathfrak{F}^{\mathbf{R}} : \mathcal{B}_p^{\mathbf{R}} \to D(X)$ in Remark 4.4. Moreover by the S-flat family of objects in $\mathcal{B}_p^{\mathbf{R}}$, their morphisms (or isomorphisms) we mean the corresponding definitions as in Definition 4.5.

4.1 The moduli stacks $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_{p,ss}}(\tilde{\tau})$ and $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}^{\mathbf{R}}_{p,ss}}(\tilde{\tau})$

Given a bounded family of τ-semistable sheaves F with fixed Hilbert polynomial P there exists an integer m such that for every sheaf F in the family, $F(m)$ is globally generated. First we construct an S-flat family of coherent sheaves F with fixed Hilbert polynomial P. To avoid confusion, here we slightly change our notation. We denote by $\mathcal{F}$ the family as a coherent $\mathcal{O}_{X\times S}$-module and by F we mean the fiber of this family over a geometric point of S. By construction, the family of coherent sheaves F appearing in a $\tilde{\tau}$-semistable rigidified object is bounded and moreover $F(n)$ is globally generated for all $n \geq m$.

Fix such n and let V be a complex vector space of dimension $d = P(n)$ given as $V = \mathrm{H}^0(F \otimes L^n)$. The line bundle L, as defined before, is the fixed polarization over X. Twisting the sheaf F by the fixed large enough n would ensure one to get a surjective morphism of coherent sheaves $V \otimes \mathcal{O}_X(-n) \to F$. One can construct a scheme parametrizing the flat quotients of $V \otimes \mathcal{O}_X(-n)$ with fixed given Hilbert polynomial. This by usual arguments provides us with Grothendieck's Quot-scheme. Here to shorten the notation we use $\mathcal{Q}$ to denote $\mathrm{Quot}_P(V \otimes \mathcal{O}_X(-n))$.

Now consider a sub-locus $\mathcal{Q}^{ss} \subset \mathcal{Q}$ which parametrizes the Giseker semistable sheaves F with fixed Hilbert polynomial P.

Definition 4.6. Define $\mathcal{P}$ over $\mathcal{Q}^{ss}$ to be the bundle whose fibers parametrize $\mathrm{H}^0(F(n))$. The fibers of the bundle $\mathcal{P}^{\oplus r}$ parametrize $\mathrm{H}^0(F(n))^{\oplus r}$. In other words the fibers of $\mathcal{P}^{\oplus r}$ parametrize the maps $\mathcal{O}_X^{\oplus r}(-n) \to F$ (which define the complexes representing the objects in $\mathcal{B}_p^{\mathbf{R}}$). Define $\mathfrak{S}_{ss}^{P,r}(\tilde{\tau}) \subset \mathcal{P}^{\oplus r}$ be given as an open subscheme of $\mathcal{P}^{\oplus r}$ whose fibers parametrize $\tilde{\tau}$-semistable objects in $\mathcal{B}_p^{\mathbf{R}}$.

Now we state (without proof) the results obtained in [15] (Section 5.2) on construction of the moduli stacks of $\tilde{\tau}$-semistable objects in $\mathcal{B}_p^{\mathbf{R}}$ and $\mathcal{B}_p$.

Theorem 4.7. *[15] (Theorem 5.7). Let $\mathfrak{S}_{ss}^{(\beta,r)}(\tilde{\tau})$ be the underlying scheme in Definition 4.6 parametrizing $\tilde{\tau}$-semistable rigidified objects of type (β, r). Let $G := \mathrm{GL}_r(\mathbb{C}) \times \mathrm{GL}(V)$. Let $\left[\frac{\mathfrak{S}_{ss}^{(\beta,r)}(\tau')}{G}\right]$ be the stack theoretic quotient of $\mathfrak{S}_{ss}^{(\beta,r)}(\tilde{\tau})$ by G. There exists an isomorphism of* *groupoids*

$$\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_p,ss}(\tilde{\tau}) \cong \left[\frac{\mathfrak{S}^{(\beta,r)}_{ss}(\tilde{\tau})}{G}\right].$$

In particular $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_p,ss}(\tilde{\tau})$ *is an Artin stack.*

Corollary 4.8. *[15] (Corollary 5.8). Apply the proof of [15] (Theorem 5.7) to* $\mathfrak{S}^{(\beta,r)}_{ss}(\tilde{\tau})$ *and* $G = \mathrm{GL}(V)$ *and obtain a natural isomorphism between* $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}^{\boldsymbol{R}}_p,ss}(\tilde{\tau})$ *and* $\left[\frac{\mathfrak{S}^{(\beta,r)}_{ss}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$.

One may use this natural isomorphism in order to obtain an alternative definition of the moduli stack of $\tilde{\tau}$-semistable rigidified objects of type (β, r) as the quotient stack $\left[\frac{\mathfrak{S}^{(\beta,r)}_{ss}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$

Corollary 4.9. *[15] (Corollary 5.9). It is true that:*

$$\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_p,ss}(\tilde{\tau}) = \left[\frac{\mathfrak{M}^{(\beta,r)}_{\mathcal{B}^{\boldsymbol{R}}_{p,ss}}(\tilde{\tau})}{\mathrm{GL}_r(\mathbb{C})}\right] \tag{4.2}$$

Proposition 4.10. *[15] (Proposition 5.10). The moduli stack,* $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}^{\boldsymbol{R}}_p,ss}(\tilde{\tau})$*, is a* $\mathrm{GL}_r(\mathbb{C})$*-torsor over* $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_p,ss}(\tilde{\tau})$*. It is true that locally in the flat topology,* $\mathfrak{M}^{(\beta,r)}_{\mathcal{B}_p,ss}(\tilde{\tau}) \cong \mathfrak{M}^{(\beta,r)}_{\mathcal{B}^{\boldsymbol{R}}_p,ss}(\tilde{\tau}) \times \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathrm{GL}_r(\mathbb{C})}\right]$*. This isomorphism does not hold true globally unless* $r = 1$*.*

5 Stack function identities in Ringel Hall algebra

Definition 5.1. (Joyce and Song) [9] (Definition 2.5). Let $\mathfrak{S}$ be a $\mathbb{C}$-stack with affine geometric stabilizers. Consider pairs $(\mathfrak{R}, \rho)$ where $\mathfrak{R}$ is a finite type algebraic $\mathbb{C}$-stack with affine geometric stabilizers and $\rho : \mathfrak{R} \to \mathfrak{S}$ is a 1-morphism. Two pairs $(\mathfrak{R}, \rho)$ and $(\acute{\mathfrak{R}}, \acute{\rho})$ are called to be equivalent if there exists a 1-morphism $\iota : \mathfrak{R} \to \acute{\mathfrak{R}}$ such that in the diagram below:

$$\begin{tikzcd} \mathfrak{R} \arrow[r, "\iota"] \arrow[d, "\rho"'] & \acute{\mathfrak{R}} \arrow[d, "\acute{\rho}"] \\ \mathfrak{S} \arrow[r, "\cong"] & \mathfrak{S} \end{tikzcd} \tag{5.1}$$

the two vertical maps are 2-isomorphic with each other. Write $[(\mathfrak{R}, \rho)]$ for the equivalence class of $(\mathfrak{R}, \rho)$. If $(\mathfrak{G}, \rho \mid_{\mathfrak{G}})$ represents a sub-pair of $(\mathfrak{R}, \rho)$ where $\mathfrak{G}$ is closed in $\mathfrak{R}$, then $(\mathfrak{G}, \rho \mid_{\mathfrak{G}})$ and $(\mathfrak{R}/\mathfrak{G}, \rho \mid_{\mathfrak{R}/\mathfrak{G}})$ are pairs of the same kind. Define $\underline{\mathrm{SF}}(\mathfrak{S})$ to be the $\mathbb{Q}$-vector space generated by equivalence classes of $[\mathfrak{R}, \rho]$ subject to the relations

$$[(\mathfrak{R}, \rho)] = [(\mathfrak{G}, \rho \mid_{\mathfrak{G}})] + [(\mathfrak{R}/\mathfrak{G}, \rho \mid_{\mathfrak{R}/\mathfrak{G}})] \tag{5.2}$$

for each subpair $(\mathfrak{G}, \rho \mid_{\mathfrak{G}})$ of $(\mathfrak{R}, \rho)$. Elements of $\underline{\mathrm{SF}}(\mathfrak{S})$ are called stack functions on $\mathfrak{S}$.

Definition 5.2. (Joyce and song) [9] (Definition 2.16.). Define the space of stack functions $\underline{\mathrm{S}}\,\overline{\mathrm{F}}(\mathfrak{S}, \chi, \mathbb{Q})$ to be the $\mathbb{Q}$-vector space generated by equivalence classes $[(\mathfrak{R}, \rho)]$ with the following relations imposed:
1. Given a sub-pair $(\mathfrak{G}, \rho\mid_{\mathfrak{G}}) \subset (\mathfrak{R}, \rho)$ we have $[(\mathfrak{R}, \rho)] = [(\mathfrak{G}, \rho\mid_{\mathfrak{G}})] + [(\mathfrak{R}/\mathfrak{G}, \rho\mid_{\mathfrak{R}/\mathfrak{G}})]$ as above.
2. Let $\mathfrak{R}$ be a $\mathbb{C}$-stack of finite type with affine geometric stabilizers and let $\mathcal{U}$ denote a quasi-projective $\mathbb{C}$-variety and $\pi_{\mathfrak{R}} : \mathfrak{R} \times U \to \mathfrak{R}$ the natural projection and $\rho : \mathfrak{R} \to \mathfrak{S}$ a 1-morphism. Then $[(\mathfrak{R} \times \mathcal{U}, \rho \circ \pi_{\mathfrak{R}})] = \chi([\mathcal{U}])[(\mathfrak{R}, \rho)]$.
3. Assume $\mathfrak{R} \cong [X/G]$ where X is a quasiprojective $\mathbb{C}$-variety and G a very special algebraic $\mathbb{C}$-group acting on X with maximal torus T^G, we have

$$[(\mathfrak{R}, \rho)] = \sum_{Q \in \mathcal{Q}(G, T^G)} F(G, T^G, Q)[([X/Q], \rho \circ \iota^Q)], \tag{5.3}$$

where the rational coefficients $F(G, T^G, Q)$ have a complicated definition explained in [5] (Section 6.2). Here $\mathcal{Q}(G, T^G)$ is the set of closed $\mathbb{C}$-subgroups Q of T^G such that $Q = T^G \cap C(G)$ and $\iota^Q : [X/Q] \to \mathfrak{R} \cong [X/G]$ is the natural projection 1-morphism, where $C(G)$ denotes the center of the group G. Similarly we can define $\overline{\mathrm{SF}}(\mathfrak{S}, \chi, \mathbb{Q})$ by restricting the 1-morphisms ρ to be representable.
Remark 5.3. There exist notions of multiplication, pullback, pushforward of stack functions in $\underline{\mathrm{S}}\,\overline{\mathrm{F}}(\mathfrak{S}, \chi, \mathbb{Q})$ and $\overline{\mathrm{SF}}(\mathfrak{S}, \chi, \mathbb{Q})$ and to save space we do not construct them here. For further discussions look at (Joyce and Song) [9] (Definitions. 2.6, 2.7) and (Theorem. 2.9).
Definition 5.4. (Joyce and Song) [9] (Definition 3.3). For $\alpha \in C(\mathcal{A})$ write $\mathfrak{M}^{\alpha}_{\mathcal{A}}$ for the substack of objects $F \in \mathcal{A}$ in class $\alpha \in K(\mathcal{A})$. Let $\mathfrak{Exact}_{\mathcal{A}}$ denote the moduli stack of short exact sequences $0 \to E_1 \to E_2 \to E_3 \to 0$ in $\mathcal{A}$. For $i = 1, 2, 3$ let $\pi_i : \mathfrak{Exact} \to \mathfrak{M}_{\mathcal{A}}$ denote the 1-morphism of Artin stacks projecting the short exact sequence $0 \to E_1 \to E_2 \to E_3 \to 0$ to E_i. Define bilinear operations $*$ on the stack function spaces $\underline{SF}$, $SF(\mathfrak{M}_{\mathcal{A}})$, $\underline{S\overline{F}}$ and $S\overline{F}(\mathfrak{M}_{\mathcal{A}}, \chi, \mathbb{Q})$ by

$$f * g = (\pi_2)_*((\pi_1 \times \pi_3)^*(f \otimes g)). \tag{5.4}$$

Definition 5.5. Let $(\tau, T, \leq)$ be a permissible stability condition on $\mathcal{A}$. Define the stack functions $\overline{\delta}^{\alpha}_{ss}(\tau) = \overline{\delta}^{\alpha}_{\mathfrak{M}^{\alpha}_{ss}}(\tau)$ in $\mathrm{SF}_{al}(\mathfrak{M}_{\mathcal{A}})$ (for definition of SF_{al} look at [9] (Definition 3.3) for $\alpha \in C(\mathcal{A})$. One thinks of $\overline{\delta}^{\alpha}_{\mathfrak{M}^{\alpha}_{ss}}(\tau)$ as the characteristic stack function of the component of the moduli stack of τ-semistable objects in $\mathcal{A}$ whose class in $K(\mathcal{A})$ is α. Define elements $\overline{\epsilon}^{\alpha}(\tau)$ in $\mathrm{SF}_{al}(\mathfrak{M}_{\mathcal{A}})$

$$\overline{\epsilon}^{\alpha}(\tau) = \sum_{\substack{n \geq 1, \alpha_1, \cdots, \alpha_n \in C(\mathcal{A}) \\ \alpha_1 + \cdots + \alpha_n = \alpha \\ \tau(\alpha_i) = \tau(\alpha) \forall i}} \frac{(-1)^{n-1}}{n} \overline{\delta}^{\alpha_1}_{ss}(\tau) * \overline{\delta}^{\alpha_2}_{ss}(\tau) * \cdots * \overline{\delta}^{\alpha_n}_{ss}(\tau), \tag{5.5}$$

where $*$ is the Ringel-Hall multiplication defined in 5.4. Here each $\overline{\delta}^{\alpha_i}_{ss}(\tau)$ denotes the characteristic stack function associated to the moduli stack of objects $F \in \mathcal{A}$ in class $\alpha_i \subset C(\mathcal{A})$ as defined in [9] (Definition 3.3).
Remark 5.6. By replacing the category $\mathcal{A}$ and the stability condition τ in Definition 5.5 with category $\mathcal{B}_p$ and $\tilde{\tau}$-stability condition respectively, we obtain the following identity over the moduli stack of $\tilde{\tau}$-semistable objects in $\mathcal{B}_p$ of type (β, r):

$$\overline{\epsilon}^{(\beta,r)}(\tilde{\tau}) = \sum_{\substack{n \geq 1, (\beta_1, d_1), \cdots, (\beta_n, d_n) \in C(\mathcal{B}_p) \\ (\beta_1, d_1) + \cdots + (\beta_n, d_n) = (\beta, r) \\ \tilde{\tau}(\beta_i, d_i) = \tilde{\tau}(\beta, d) \forall i}} \frac{(-1)^{n-1}}{n} \overline{\delta}^{(\beta_1,d_1)}_{ss}(\tau) * \overline{\delta}^{(\beta_2,d_2)}_{ss}(\tau) * \cdots * \overline{\delta}^{(\beta_n,d_n)}_{ss}(\tau). \tag{5.6}$$

Note that Equation (5.6) will be the key identity throughout this article. In the remaining sections we intend to compute the invariants associated to the moduli stack of $\tilde{\tau}$-semistable objects of type (β, r) in $\mathcal{B}_p$ for when $r = 2$ by directly computing the Ringel Hall product appearing on the right hand side of (5.6).

Now we apply the Lie algebra morphism $\tilde{\Psi}^{\mathcal{B}_p}$ defined in [9] (Section 13.4) to the left side of Equation (5.6) and define the invariant assciated to the moduli stack of objects of type (β, r) in $\mathcal{B}_p$:

Definition 5.7. Define the invariant $\mathbf{B}_p^{ss}(X, \beta, 2, \tilde{\tau})$ associated to $\tilde{\tau}$-semistable objects of type $(\beta, 2)$ in $\mathcal{B}_p$ by

$$\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})) = \mathbf{B}_p^{ss}(X, \beta, 2, \tilde{\tau}) \cdot \tilde{\lambda}^{(\beta,2)},$$

where $\tilde{\Psi}^{\mathcal{B}_p}$ is given by the Lie algebra morphism defined in [9] (Section 13.4).

6 Direct computation of invariants in an example

In this section we introduce a direct approach to the calculation of the invariants of objects in $\mathcal{B}_p$ via carrying out an example. Our calculation exploits the properties of stack functions defined by Joyce [5], [6], [4], [7], [8] but it does not involve any wallcrossing computation. We will show that our results in this section verify the results obtained in [15] (Equation 12.32) through an example.

Example 6.1. Computation of $\mathbf{B}_p^{ss}(X, [\mathbb{P}^1], 2, \tilde{\tau})$ where X is given by total space of $\mathcal{O}_{\mathbb{P}^1}^{\oplus 2}(-1) \to \mathbb{P}^1$.

We compute the invariant of $\tilde{\tau}$-semistable objects $(F, \mathbb{C}^2, \phi_{\mathbb{C}^2})$ of type $([\mathbb{P}^1], 2)$ in $\mathcal{B}_p$. Note that In this case F has rank 1 and $p(n) = n + \chi(F)$. Assume $\chi(F) = r$. In this case by computations in [11] and [3] the only semistable sheaf, F, with $\mathrm{ch}_2(F) = [\mathbb{P}^1]$ is given by $\mathcal{O}_{\mathbb{P}^1}(r-1)$ which is a stable sheaf. First we give description of $\mathfrak{M}_{ss,\mathcal{B}_p}^{(2,[\mathbb{P}^1])}(\tilde{\tau})$. By definition an object of type $([\mathbb{P}^1, 2])$ in $\mathcal{B}_p$ is identified by a complex $\mathcal{O}_X(-n)^{\oplus 2} \to \iota_* \mathcal{O}_{\mathbb{P}^1}(r-1)$ where $\iota : \mathbb{P}^1 \hookrightarrow X$ (from now on we suppress ι_* in our notation). By the constructions in Section 4.1 the parameter scheme of $\tilde{\tau}$-semistable objects is obtained by choosing two sections (s_1, s_2) such that $s_i \in \mathrm{H}^0((\mathcal{O}_{\mathbb{P}^1}(n+r-1))$ for $i = 1, 2$. More over since $\mathcal{O}_{\mathbb{P}^1}(r-1)$ is a stable sheaf, its stabilizer is given by $\mathbb{G}_m$.

An important point to note is that given a $\tilde{\tau}$-semistable object $(F, \mathbb{C}^2, \phi_{\mathbb{C}^2})$ one is always able to obtain a an exact sequence of the form

$$0 \to (F, \mathbb{C}, \phi_{\mathbb{C}}) \to (F, \mathbb{C}^2, \phi_{\mathbb{C}^2}) \to (0, \mathbb{C}, 0) \to 0, \tag{6.1}$$

for every object in the moduli stack and since $\tilde{\tau}(F, \mathbb{C}^2, \phi_{\mathbb{C}}) = 1 \leq \tilde{\tau}(0, \mathbb{C}, 0) = 1$ then one concludes that all the objects parametrized by the moduli stack are given by extensions of rank 1 $\tilde{\tau}$-stable objects and hence all objects are $\tilde{\tau}$-strictly-semistable. Moreover, note that giving a $\tilde{\tau}$-semistable object of the form $\mathcal{O}_X(-n)^{\oplus 2} \xrightarrow{(s_1,s_2)} F$ is equivalent to requiring the condition that $(s_1, s_2) \neq (0, 0)$, since other wise one may be able to obtain a an exact sequence:

$$0 \to (\mathbb{C}^2, 0, 0) \to (\mathbb{C}^2, F, 0) \to (0, F, 0) \to 0$$

such that $\tilde{\tau}(\mathbb{C}^2, 0, 0) = 1 > \tilde{\tau}(0, F, 0) = 0$, hence $(\mathbb{C}^2, 0, 0)$ weakly destabilizes $(\mathbb{C}^2, F, 0)$ and one obtains a contradiction. Now use Theorem 4.7 and obtain

$$\mathfrak{M}_{ss,\mathcal{B}_p}^{(2,[\mathbb{P}^1])}(\tilde{\tau}) = \left[\frac{(\mathrm{H}^0((\mathcal{O}_{\mathbb{P}^1}(n+r-1)))^{\oplus 2} \backslash \{0\} / \mathbb{G}_m}{\mathrm{GL}_2(\mathbb{C})}\right] \cong \left[\frac{\mathbb{P}(\mathrm{H}^0((\mathcal{O}_{\mathbb{P}^1}(n+r-1))^{\oplus 2})}{\mathrm{GL}_2(\mathbb{C})}\right]. \tag{6.2}$$

Now we need to compute the element of the Hall algebra $\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})$. By applying Definition 5.6 to $\mathfrak{M}^{([\mathbb{P}^1],2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$ we obtain:

$$\bar{\epsilon}^{(\beta,2)}(\tilde{\tau}) = \bar{\delta}^{([\mathcal{P}^1],2)}_{ss}(\tilde{\tau}) - \frac{1}{2}\sum_{\beta_k+\beta_l=[\mathbb{P}^1]} \bar{\delta}^{(\beta_k,1)}_{s}(\tilde{\tau}) * \bar{\delta}^{(\beta_l,1)}_{s}(\tilde{\tau}). \tag{6.3}$$

Now we use a stratification strategy in order to decompose $\mathfrak{M}^{([\mathbb{P}^1],2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$ into a disjoint union of strata as follows: Given the fact that the objects in the moduli stack under study are of type $([\mathbb{P}^1],2)$ one would immediately see that the only possible decomposition for a $\tilde{\tau}$-semistable object of type $([\mathbb{P}^1],2)$ is given by decomposition of its class as $([\mathbb{P}^1],2) = ([\mathbb{P}^1],1) + (0,1)$. This means that a strictly $\tilde{\tau}$-semistable object of type $([\mathbb{P}^1],2)$ is either given by (split or non-split) extensions of object of type $([\mathbb{P}^1],1)$ by objects of type $(0,1)$ or it is given by the extensions with reversed order, i.e the extensions of objects of type $(0,1)$ by objects of type $([\mathbb{P}^1],1)$. Our stratification technique involves a study of the parametrizing moduli stacks for these objects depending on what extensions are used to produce the objects. We decompose $\mathfrak{M}^{([\mathbb{P}^1],2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$ into a disjoint union of split and non-split strata.

Definition 6.2. Define $\mathfrak{M}^{([\mathbb{P}^1],2)}_{sp,\mathcal{B}_p}(\tilde{\tau}) \subset \mathfrak{M}^{([\mathbb{P}^1],2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$ to be the locally closed stratum over which an object of type $([\mathbb{P}^1],2)$ is given by split extensions involving objects of type $([\mathbb{P}^1],1)$ and $(0,1)$. Define $\mathfrak{M}^{([\mathbb{P}^1],2)}_{nsp,\mathcal{B}_p}(\tilde{\tau}) \subset \mathfrak{M}^{([\mathbb{P}^1],2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$ to be a locally closed stratum over which an object of type $([\mathbb{P}^1],2)$ is given by non-split extensions involving objects of type $([\mathbb{P}^1],1)$ and $(0,1)$.

Now we study the structure of each stratum separately.

6.1 Stacky structure of $\mathfrak{M}^{([\mathbb{P}^1],2)}_{sp,\mathcal{B}_p}(\tilde{\tau})$

It is easy to see that any $\tilde{\tau}$-semistable objects of type $([\mathbb{P}^1],2)$ given as

$$\mathcal{O}_X(-n)^{\oplus 2} \to \mathcal{O}_{\mathbb{P}^1}(r-1) \cong \big(\mathcal{O}_X(-n)^{\oplus 2} \to \mathcal{O}_{\mathbb{P}^1}(r-1)\big) \oplus (\mathcal{O}_X(-n) \to 0)$$

has the property that the sections s_1, s_2 for this object are linearly dependent on one another. Hence by discussion Definition 4.6 the underlying parameter scheme of $\tilde{\tau}$-semistable objects of this given form is given by chooing a nonzero section of $\mathcal{O}_{\mathbb{P}^1}(n+r-1)$ in other words we obtain $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1))\backslash\{0\}$. Now we need to take the quotient of this space by the stabilizer group of points. We know that the condition required for a $\tilde{\tau}$-semistable object $\mathcal{O}_X(-n)^{\oplus 2} \xrightarrow{(s_1,s_2)} F$ to be given by split extensions of rank 1 objects is that s_1 and s_2 are linearly dependent on one another. Now pick such an object given by $\mathcal{O}_X(-n)^{\oplus 2} \xrightarrow{(s_1,0)} F$. The automorphisms of this object are given by the group which makes the following diagram commutative:

$$\begin{tikzcd}
\mathcal{O}_X(-n)^{\oplus 2} \arrow[r, "{(s_1,0)}"] \arrow[d, "\cong"'] & \mathcal{O}_{\mathbb{P}^1}(r-1) \arrow[d, "\cong"] \\
\mathcal{O}_X(-n)^{\oplus 2} \arrow[r, "{(s_1,0)}"] & \mathcal{O}_{\mathbb{P}^1}(r-1)
\end{tikzcd}$$

Hence it is seen that the left vertical map needs to be given by a subgroup of $\mathrm{GL}_2(\mathbb{C})$ which preserves s_1, i.e the Borel subgroup of $\mathrm{GL}_2(\mathbb{C})$ whose elements are given by 2×2 upper triangular matrices $\begin{pmatrix} k_1 & k_2 \\ 0 & k_3 \end{pmatrix}$ where $k_1, k_3 \in \mathbb{G}_m$ and $k_2 \in \mathbb{A}^1$. Having fixed one of the automorphisms via fixing k_1, k_2, k_3, it is seen that by the commutativity of the square diagram the right vertical map needs to be given by multiplication by k_1 which is an element of $\mathbb{G}_m$. Note that one needs to take the quotient of the parameter scheme by all isomorphisms between any two objects in the split stratum, not just the automorphisms of one fixed representative. In general for an object to live in the split stratum one requires the sections (s_1, s_2) to be given by $(s_1, a\cdot s_1)$. We observed that fixing a representative for a split object of rank 2 (such as fixing $(s_1, s_2) = (s_1, 0)$ as above) would tell us that its automorphisms are given by $\mathbb{G}_m^2 \rtimes \mathbb{A}^1$. Hence taking into account all possible representatives implies that the stabilizer group of objects in spit stratum is given by $\mathbb{G}_m^2 \rtimes \mathbb{A}^1 \times \mathbb{G}_m$. Hence we obtain

$$\mathfrak{M}^{([\mathbb{P}^1],2)}_{sp,\mathcal{B}_p}(\tilde{\tau}) = \left[\frac{\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1))\backslash\{0\}}{\mathbb{G}_m^2 \rtimes \mathbb{A}^1 \times \mathbb{G}_m}\right] = \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2 \rtimes \mathbb{A}^1}\right]. \tag{6.4}$$

6.2 Stacky structure of $\mathfrak{M}^{([\mathbb{P}^1],2)}_{nsp,\mathcal{B}_p}(\tilde{\tau})$

In this case all the objects in $\mathfrak{M}^{([\mathbb{P}^1],2)}_{nsp,\mathcal{B}_p}(\tilde{\tau})$ are given by non-split extensions of the form:

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{O}_X(-n) \arrow[r] \arrow[d, "s_1"] & \mathcal{O}_X^{\oplus 2}(-n) \arrow[r] \arrow[d, "{(s_1,s_2)}"] & \mathcal{O}_X(-n) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \mathcal{O}_{\mathbb{P}^1}(r-1) \arrow[r, "\cong"] & F \arrow[r] & 0 \arrow[r] & 0,
\end{tikzcd} \tag{6.5}$$

Note that switching the place of $\mathcal{O}_{\mathbb{P}^1}(r-1)$ and 0 in the bottom row of diagram (6.5) would produce a split extension. Now in order to obtain non-split extensions one needs to choose two sections s_1, s_2 such that s_1 and s_2 are linearly independent. The set of all linearly independent choices of s_1 and s_2 spans a two dimensional subspace of $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1))$ which is given by the Grassmanian:

$$\mathrm{G}(2, n+r).$$

Now we need to take the quotient of this scheme by the stabilizer group of points in the stratum. We know that the condition required for a $\tilde{\tau}$-semistable object $\mathcal{O}_X(-n)^{\oplus 2} \xrightarrow{(s_1,s_2)} F$ to be given by nonsplit extensions of rank 1 objects is that s_1 and s_2 are linearly independent. Now pick such an object given by $\mathcal{O}_X(-n)^{\oplus 2} \xrightarrow{(s_1,s_2)} F$. The automorphisms of this object are given by the group which makes the following diagram commutative:

$$\begin{tikzcd}
\mathcal{O}_X(-n)^{\oplus 2} \arrow[r, "{(s_1,s_2)}"] \arrow[d, "\cong"'] & \mathcal{O}_{\mathbb{P}^1}(r-1) \arrow[d, "\cong"] \\
\mathcal{O}_X(-n)^{\oplus 2} \arrow[r, "{(s_1,s_2)}"] & \mathcal{O}_{\mathbb{P}^1}(r-1)
\end{tikzcd}$$

Hence it is seen that the left vertical map needs to be given by a subgroup of $\mathrm{GL}_2(\mathbb{C})$ whose elements are given by 2×2 diagonal matrices of the form $\begin{pmatrix} k_1 & 0 \\ 0 & k_1 \end{pmatrix}$ where $k_1 \in \mathbb{G}_m$. Having fixed one of the automorphisms via fixing k_1, it is seen that by the commutativity of the square diagram the right vertical map needs to be given by multiplication by k_1 which is an element of $\mathbb{G}_m$. Hence we obtain

$$\mathfrak{M}^{([\mathbb{P}^1],2)}_{nsp,\mathcal{B}_p}(\tilde{\tau}) = \left[\frac{\mathrm{G}(2,n+r)}{\mathbb{G}_m}\right]. \tag{6.6}$$

Now we compute $\sum_{\beta_k+\beta_l=[\mathbb{P}^1]} \bar{\delta}^{(\beta_k,1)}_s(\tilde{\tau}) * \bar{\delta}^{(\beta_l,1)}_s(\tilde{\tau})$ appearing on the right hand side of (6.3). We use the fact that

$$\sum_{\beta_k+\beta_l=[\mathbb{P}^1]} \bar{\delta}^{(\beta_k,1)}_s(\tilde{\tau}) * \bar{\delta}^{(\beta_l,1)}_s(\tilde{\tau}) = \bar{\delta}^{([\mathbb{P}^1],1)}_s(\tilde{\tau}) * \bar{\delta}^{(0,1)}_s(\tilde{\tau}) + \bar{\delta}^{(0,1)}_s(\tilde{\tau}) * \bar{\delta}^{([\mathbb{P}^1],1)}_s(\tilde{\tau}) \tag{6.7}$$

and compute each term on the right hand side of (6.7) separately.

Remark 6.3. As we described above there exists an action of $\mathrm{GL}_2(\mathbb{C})$ on $\mathbf{S} := \mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1))^{\oplus 2}))$. This action induces an action of the corresponding Lie algebra on the tangent space of $\mathbf{S}$ given by the map:

$$\mathcal{O}_{\mathbf{S}} \otimes \mathfrak{K} \to T_{\mathbf{S}}, \tag{6.8}$$

where $\mathfrak{K}$ denotes the Lie algebra associated to the group $\mathrm{GL}_2(\mathbb{C})$. The dimension of the automorphism group of objects representing the elements of $\mathbf{S}$ is given by the dimension of the stabilizer (in $\mathrm{GL}_2(\mathbb{C})$) group of these elements, which itself is given by the dimension of the kernel of the map in (8.8). On the other hand, the dimension of the kernel of the map in (8.8) is an upper-semicontinious function. Therefore by the usual arguments, we obtain a stratification of $\mathbf{S}$ which induces a stratification of $\left[\frac{\mathbf{S}}{\mathrm{GL}_2(\mathbb{C})}\right]$ into locally closed strata such that over each stratum the dimension of the stabilizer group is constant as we vary over points inside that stratum. Hence in Definition 6.2 we stated without proof that the defined strata are locally closed in $\mathfrak{M}^{([\mathbb{P}^1],2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$.

6.3 Computation of $\bar{\delta}^{([\mathbb{P}^1],1)}_s(\tilde{\tau}) * \bar{\delta}^{(0,1)}_s(\tilde{\tau})$

Given $\bar{\delta}^{([\mathbb{P}^1],1)}_s(\tilde{\tau}) = (\left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right], \rho_1)$ and $\bar{\delta}^{(0,1)}_s(\tilde{\tau}) = (\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right], \rho_2)$ consider the diagram:

$$\begin{tikzcd}
\mathcal{Z}'_{12} \arrow[r, "\Phi"] \arrow[d] & \mathfrak{Exact}_{\mathcal{B}_p} \arrow[r, "\pi_2"] \arrow[d, "\pi_1 \times \pi_3"] & \left[\frac{\mathbb{P}(\mathrm{H}^0((\mathcal{O}_{\mathbb{P}^1}(n+r-1))^{\oplus 2})}{\mathrm{GL}_2(\mathbb{C})}\right] \\
\left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right] \times \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \arrow[r, "\rho_1 \times \rho_2"] & \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) \times \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) &
\end{tikzcd} \tag{6.9}$$

Here $\mathcal{Z}'_{12}$ is given by the scheme parametrizing the set of commutative diagrams:

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{O}_X(-n) \arrow[r] \arrow[d, "s_1"] & \mathcal{O}_X^{\oplus 2}(-n) \arrow[r] \arrow[d, "(s_1, s_2)"] & \mathcal{O}_X(-n) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & \mathcal{O}_{\mathbb{P}^1}(r-1) \arrow[r, "\cong"] & F \arrow[r] & 0 \arrow[r] & 0,
\end{tikzcd} \tag{6.10}$$

Note that the extensions in (6.10) have the possibility of being split or non-split. Hence we consider each case separately and introduce a new notation:

Definition 6.4. Let $[\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau})]_{sp}$ denote the stratum of $(\pi_2 \circ \Phi)_* \mathcal{Z}'_{12}$ over which the points are represented by split extensions given by coomutative diagram in (6.10).
Definition 6.5. Let $[\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau})]_{nsp}$ denote the stratum of $(\pi_2 \circ \Phi)_* \mathcal{Z}'_{12}$ over which the points are represented by non-split extensions given by coomutative diagram in (6.10).

Now we compute $[\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau})]_{sp}$. This amounts to choosing sections s_1, s_2 so that s_1 and s_2 are linearly depending on one another. The scheme parametrizing the nonzero sections s_1 is given by $\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))$. Now we take the quotient of this scheme by the stabilizer group of points. Similar to arguments in [15] (Lemma 12.1) given any point in $\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))$ represented by the extension in diagram (6.10) its stabilizer group is given by the semi-direct product $\mathbb{G}_m^2 \rtimes \mathrm{Hom}(E_3, E_1)$ where each factor of $\mathbb{G}_m$ amounts to the stabilizer group of objects given as $E_3 := \mathcal{O}_X(-n) \to 0$ and $E_1 := \mathcal{O}_X(-n) \to \mathcal{O}_{\mathbb{P}^1}(r-1)$ respectively. Note that the extra factor of $\mathbb{A}^1$ will not appear as a part of the stabilizer group since by the given description of E_1 and E_3 we know that $\mathrm{Hom}(E_3, E_1) = 0$ for every such E_1 and E_3. We obtain the following conclusion:

$$[\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau})]_{sp} = \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2}\right]. \tag{6.11}$$

Now we compute $[\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau})]_{nsp}$. This amounts to choosing s_1, s_2 so that s_1 and s_2 are linearly independent and the extension in diagram (6.10) becomes non-split . Note that for any fixed value of s_1 one has $\mathbb{P}^1$ worth of choices for s_2. Now we need to consider all possible choices of s_1 and in doing so we require the pair s_1, s_2 to remain linearly independent. This gives the flag variety $\mathrm{F}(1, 2, n+r)$. Hence we obtain:

$$[\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau})]_{nsp} = \left[\frac{\mathrm{F}(1, 2, n+r)}{\mathbb{G}_m}\right]. \tag{6.12}$$

Note that the factor of $\mathbb{G}_m$ in the denominator of (6.12) is due to the fact that we have used one of the $\mathbb{G}_m$ factors in projectivising the bundle of s_2-choices over the Grassmanian. We finish this section by summarizing our computation. By (6.11) and (6.12) one obtains:

$$\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau}) = \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2}\right] + \left[\frac{\mathrm{F}(1, 2, n+r)}{\mathbb{G}_m}\right] \tag{6.13}$$

6.4 Computation of $\bar{\delta}_s^{(0,1)}(\tilde{\tau}) * \bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau})$

Now change the order of $\bar{\delta}_s^{(0,1)}(\tilde{\tau})$ and $\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau})$ and obtain a diagram

$$\begin{tikzcd}
\mathcal{Z}'_{21} \arrow[r, "\Phi"] \arrow[d] & \mathfrak{Exact}_{\mathcal{B}_p} \arrow[r, "\pi_2"] \arrow[d, "\pi_1 \times \pi_3"] & \left[\frac{\mathbb{P}(\mathrm{H}^0((\mathcal{O}_{\mathbb{P}^1}(n+r-1))^{\oplus 2})}{\mathrm{GL}_2(\mathbb{C})}\right] \\
\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \times \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right] \arrow[r, "\rho_2 \times \rho_1"] & \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) \times \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) &
\end{tikzcd} \tag{6.14}$$

Here $\mathcal{Z}'_{21}$ is given by the scheme parametrizing the set of commutative diagrams:

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{O}_X(-n) \arrow[r] \arrow[d, "0"] & \mathcal{O}_X^{\oplus 2}(-n) \arrow[r] \arrow[d, "(0,s_2)"] & \mathcal{O}_X(-n) \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & 0 \arrow[r, "\cong"] & F \arrow[r] & \mathcal{O}_{\mathbb{P}^1}(r-1) \arrow[r] & 0,
\end{tikzcd} \tag{6.15}$$

Note that the computation in this case is easier since the only possible extensions of the form given in (6.15) are the split extensions. The computation in this case is similar to computation in (6.11) except that one needs to take into account that over any point represented by an extension (as in diagram (6.15)) of $E_1 := \mathcal{O}_X(-n) \to 0$ and $E_3 := \mathcal{O}_X(-n) \to \mathcal{O}_{\mathbb{P}^1}(r-1)$ we have $\mathrm{Hom}(E_3, E_1) \cong \mathbb{A}^1$. Hence by similar discussions we obtain :

$$\bar{\delta}_s^{([\mathbb{P}^1],1)}(\tilde{\tau}) * \bar{\delta}_s^{(0,1)}(\tilde{\tau}) = \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2 \rtimes \mathbb{A}^1}\right]. \tag{6.16}$$

6.5 Computation of $\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})$

By (6.3), (6.4), (6.6), (6.7), (6.13) and (6.16) we obtain:

$$\begin{aligned}
\bar{\epsilon}^{(\beta,2)}(\tilde{\tau}) = & \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2 \rtimes \mathbb{A}^1}\right] + \left[\frac{\mathrm{G}(2,n+r)}{\mathbb{G}_m}\right] \\
& - \frac{1}{2} \cdot \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2}\right] - \frac{1}{2} \cdot \left[\frac{\mathrm{F}(1,2,n+r)}{\mathbb{G}_m}\right] \\
& - \frac{1}{2} \cdot \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2 \rtimes \mathbb{A}^1}\right].
\end{aligned} \tag{6.17}$$

Now use the decomposition used by Joyce and Song in [9] (page 158) and write

$$\begin{aligned}
& \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2 \rtimes \mathbb{A}^1}\right] = \\
& F(G, \mathbb{G}_m^2, \mathbb{G}_m^2) \cdot \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2}\right] + F(G, \mathbb{G}_m^2, \mathbb{G}_m) \cdot \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right],
\end{aligned} \tag{6.18}$$

where $F(G, \mathbb{G}_m^2, \mathbb{G}_m^2) = 1$ and $F(G, \mathbb{G}_m^2, \mathbb{G}_m) = -1$. Equation (6.17) simplifies as follows:

$$\begin{aligned}
\bar{\epsilon}^{(\beta,2)}(\tilde{\tau}) = &\cancel{\left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2}\right]} - \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right] \\
&+ \left[\frac{\mathrm{G}(2, n+r)}{\mathbb{G}_m}\right] - \frac{1}{2} \cdot \cancel{\left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2}\right]} \\
&\frac{1}{2} \cdot \left[\frac{\mathrm{F}(1, 2, n+r)}{\mathbb{G}_m}\right] - \frac{1}{2} \cdot \cancel{\left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m^2}\right]} \\
&+ \frac{1}{2} \cdot \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right] \qquad (6.19)
\end{aligned}$$

Now use Definition 9.1 and write:

$$\left[\frac{\mathrm{G}(2, n+r)}{\mathbb{G}_m}\right] = \chi(\mathrm{G}(2, n+r)) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \qquad (6.20)$$

and

$$\begin{aligned}
&\left[\frac{\mathrm{F}(1, 2, n+r)}{\mathbb{G}_m}\right] = \\
&= \chi(\mathrm{F}(1, 2, n+r)) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \\
&= \chi(\mathbb{P}^1) \cdot \chi(\mathrm{G}(2, n+r)) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \\
&= 2 \cdot \chi(\mathrm{G}(2, n+r)) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \qquad (6.21)
\end{aligned}$$

where the equality in the third line is due the fact that the topological Euler characteristic of a vector bundle over a base variety is equal to the Euler characteristic of its fibers times the Euler characteristic of the base. By (6.17) and (6.21) we obtain:

$$\begin{aligned}
&- \frac{1}{2} \cdot \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right] + \chi(\mathrm{G}(2, n+r)) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \\
&- 2 \cdot \frac{1}{2} \cdot \chi(\mathrm{G}(2, n+r)) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \\
&= -\frac{1}{2} \cdot \left[\frac{\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))}{\mathbb{G}_m}\right] = -\frac{1}{2}\chi(\mathbb{P}(\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(n+r-1)))) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \\
&= -\frac{1}{2}(n+r) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]. \qquad (6.22)
\end{aligned}$$

6.6 Computation of the invariant

Now apply the Lie algebra morphism $\tilde{\Psi}^{\mathcal{B}_p}$ [9] (Section 13.4) to $\bar{\epsilon}^{([\mathbb{P}^1],2)}(\tilde{\tau})$. By definition:

$$\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{([\mathbb{P}^1],2)}(\tilde{\tau})) = \chi^{na}\left(-\frac{1}{2}(n+r) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right], (\mu \circ i_2)^* \nu_{\mathfrak{M}^{(0,2)}_{\mathcal{B}_p}}\right)\tilde{\lambda}^{([\mathbb{P}^1],2)}. \qquad (6.23)$$

Note that by Equation (6.2) $\mathfrak{M}^{(2,[\mathbb{P}^1])}_{ss,\mathcal{B}_p}(\tilde{\tau}) = \left[\frac{\mathbb{P}(\mathrm{H}^0((\mathcal{O}_{\mathbb{P}^1}(n+r-1))^{\oplus 2})}{\mathrm{GL}_2(\mathbb{C})}\right]$ and hence $\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]$ has relative dimension $-1-(2n+2r-5) = 4-2n-2r$ over $\mathfrak{M}^{(2,[\mathbb{P}^1])}_{ss,\mathcal{B}_p}(\tilde{\tau})$. Moreover, $\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]$ is given by a single point with Behrend's multiplicity -1 and

$$(\mu \circ i_2)^* \nu_{\mathfrak{M}^{(0,2)}_{\mathcal{B}_p}})\tilde{\lambda}^{([\mathbb{P}^1],2)} = (-1)^{4-2n-2r} \cdot \nu_{\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]} = \nu_{\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]},$$

therefore:

$$\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{([\mathbb{P}^1],2)}(\tau^\bullet)) = \chi^{na}\left(-\frac{1}{2}(n+r)\cdot\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right], \nu_{\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]}\right)\tilde{\lambda}^{([\mathbb{P}^1],2)} = (-1)^1 \cdot \frac{-1}{2}(n+r)\cdot\tilde{\lambda}^{([\mathbb{P}^1],2)}. \tag{6.24}$$

Finally by Definition 5.7 we obtain:

$$\mathbf{B}^{ss}_p(X,\beta,2,\tilde{\tau}) = \frac{1}{2}(n+r). \tag{6.25}$$

Note that it is easily seen that substituting $([\mathbb{P}^1],2)$ for $(\beta,2)$ in [15] (Equation 12.32) would give the same answer as in (6.25).

7 An approach to direct calculations in general cases

In section 6 we introduced our strategy of direct computation in an example where $r=2$ and β was given by the irreducible class $[\mathbb{P}^1]$. Our strategy involved computing the weighted Euler characteristic of the right hand side of Equation (6.3). After giving the stacky structure of the moduli space of objects of type $([\mathbb{P}^1],2)$ (Section 6.2) first we computed the second summand on the right hand side of Equation (6.3) (sections 6.3 and 6.4). Then we considered a stratification of the original moduli stack so that the first summand on the right hand side of Equation (6.3) is written as a sum of the characteristic stack functions of the strata in a way so that after suitable cancellations (Equation (6.19)) the right hand side of Equation (6.3) is given as a sum of stack functions supported over virtual indecomposables. Finally by applying the Lie algebra morphism defined in [9] (Section 13.4) to both sides of Equation (6.3) we obtained Equation (6.25). In this section we adapt the same computational strategy to the case where $r=2$ and β has the possibility of being given by an irreducible class:

Assumption 7.1. Throughout this section we assume that a $\tilde{\tau}$-semistable object $(F,V,\phi_V)\in\mathcal{B}_p$ of type $(\beta,2)$ has the property that β is either indecomposable or it satisfies the condition that if $\beta=\beta_k+\beta_l$ (i.e if β is decomposable) then $\beta_k=[F_k]$ and $\beta_l=[F_l]$ such that F_k and F_l are Giseker **stable** sheaves with fixed reduced Hilbert polynomial p. In other words β can not be decomposed into smaller classes whose associated sheaves are not Giseker stable.

Lemma 7.2. *Let $\mathfrak{S}^{(\beta_k,1)}_s(\tilde{\tau})$ and $\mathfrak{S}^{(\beta_l,1)}_s(\tilde{\tau})$ (for some β_k and β_l) denote the underlying parametrizing schemes as in Definition 4.6 (for the case where r is set to be equal to 1 and $\beta=\beta_k$ and $\beta=\beta_l$ respectively). Moreover, assume that β_k and β_l satisfy the condition in Assumption 7.1. Given a tuple of stable objects, $(E_1,E_3)\in\mathfrak{S}^{(\beta_k,1)}_s(\tilde{\tau})\times\mathfrak{S}^{(\beta_l,1)}_s(\tilde{\tau})$, the following is true:*

$$\mathrm{Hom}(E_3,E_1)=\mathbb{A}^1,$$

if $E_1 \cong E_3$ *and*

$$\mathrm{Hom}(E_3, E_1) = 0,$$

if $E_1 \ncong E_3$.

Proof. . Fix $E_1 := [\mathcal{O}_X(-n) \to F_1] \in \mathfrak{S}_s^{(\beta_k,1)}(\tilde{\tau})$ and $E_3 := [\mathcal{O}_X(-n) \to F_3] \in \mathfrak{S}_s^{(\beta_l,1)}(\tilde{\tau})$. Consider a map $\psi : E_1 \to E_3$. By definition the morphism between the $\tilde{\tau}$-stable objects E_1 and E_3 is defined by a morphism $\psi_F : F_1 \to F_3$ which makes the following diagram commutative:

$$\begin{tikzcd}
O_X(-n) \arrow[r] \arrow[d, "id_{\mathcal{O}_X}"'] & F_1 \arrow[d, "\psi_F"] \\
O_X(-n) \arrow[r] & F_3
\end{tikzcd} \tag{7.1}$$

By assumption F_1 and F_3 are given as stable sheaves with fixed reduced Hilbert polynomial p. Hence any nontrivial sheaf homomorphism from F_1 to F_3 is an isomorphism. Moreover by simplicity of stable sheaves any such nontrivial isomorphism is identified with $\mathbb{A}^1$. Now use the commutativity of the diagram in (7.1). ☐

Lemma 7.3. *Fix* β_k *and* β_l *such that* $\beta_k + \beta_l = \beta$. *Consider a* $\tilde{\tau}$*-semistable object* E_2 *given as an element of* $\mathfrak{S}_{ss}^{(\beta,2)}(\tilde{\tau})$ *which fits into a non-split extension*

$$0 \to E_1 \to E_2 \to E_3 \to 0,$$

where E_1 *and* E_3 *are* $\tilde{\tau}$*-stable objects with classes* $(\beta_k, 1)$ *and* $(\beta_l, 1)$ *respectively (i.e given by elements of* $\mathfrak{S}_s^{(\beta_k,1)}(\tilde{\tau})$ *and* $\mathfrak{S}_s^{(\beta_l,1)}(\tilde{\tau})$ *respectively). Assume that* β, β_k *and* β_l *satisfy the condition in Assumption 7.1. Suppose* E_1' *and* E_3' *are* $\tilde{\tau}$*-stable objects with classes* $(\beta_{k'}, 1)$ *and* $(\beta_{l'}, 1)$ *respectively where* $\beta_{k'} + \beta_{l'} = \beta$ *and neither* β_k *nor* β_l *are equal to* $\beta_{k'}$ *or* $\beta_{l'}$. *Assume that* β, β_k' *and* β_l' *also satisfy the condition in Assumption 7.1. Then* E_2 *can not be given as an extension*

$$0 \to E_1' \to E_2 \to E_3' \to 0.$$

Proof. . If E_2 is given by both extensions then we obtain a map between the two sequences:

$$\begin{tikzcd}
0 \arrow[r] & E_1 \arrow[r, "\iota"] & E_2 \arrow[r] \arrow[d, "\cong"] & E_3 \arrow[r] & 0 \\
0 \arrow[r] & E_1' \arrow[r] & E_2 \arrow[r, "p"'] & E_3' \arrow[r] & 0
\end{tikzcd} . \tag{7.2}$$

Hence we obtain a map $p \circ \iota : E_1 \to E_3'$. Since by assumption $\beta_k \neq \beta_{k'}$ (this means $\beta_l \neq \beta_{l'}$ because $\beta_k + \beta_l = \beta_{k'} + \beta_{l'} = \beta$), by Lemma 7.2 we conclude that $p \circ \iota$ is the zero map. Hence $p \circ \iota$ factors through the map $\iota' \circ g$ in the following diagram

$$\begin{tikzcd}
0 \arrow[r] & E_1 \arrow[r, "\iota"] \arrow[d, "g"] & E_2 \arrow[r] \arrow[d, "\cong"] & E_3 \arrow[r] & 0 \\
0 \arrow[r] & E_1' \arrow[r, "\iota'"] & E_2 \arrow[r, "p"'] & E_3' \arrow[r] & 0
\end{tikzcd} . \tag{7.3}$$

Since $E_1 \ncong E_1'$, by Lemma 7.2, g is the zero map. By considering the left commutative square in (8.4) we obtain a contradiction since the image of E_1 in E_2 can not always be zero. □

Lemma 7.4. *Fix β_k and β_l such that $\beta_k + \beta_l = \beta$. Consider a $\tilde{\tau}$-semisable object E_2 with class $(\beta, 2)$ which fits into a non-split extension*

$$0 \to E_1 \to E_2 \to E_3 \to 0,$$

where E_1 and E_3 are $\tilde{\tau}$-stable objects with classes $(\beta_k, 1)$ and $(\beta_l, 1)$ respectively. Assume that β, β_k and β_l satisfy the condition in Assumption 7.1. The object E_2 can not be given as an extension

$$0 \to E_1' \to E_2 \to E_3' \to 0,$$

where E_1' is $\tilde{\tau}$-stable with class $(\beta_l, 1)$ and E_3' is $\tilde{\tau}$-stable with class $(\beta_k, 1)$.

Proof. . We prove by contradiction. Assume E_2 fits in both exact sequences. We obtain a map between the two sequences

$$\begin{tikzcd}
0 \arrow[r] & E_1 \arrow[r, "\iota"] & E_2 \arrow[r] \arrow[d, "\cong"] & E_3 \arrow[r] & 0 \\
0 \arrow[r] & E_1' \arrow[r] & E_2 \arrow[r, "p"'] & E_3' \arrow[r] & 0
\end{tikzcd} . \tag{7.4}$$

Similar to before, we obtain a map $p \circ \iota : E_1 \to E_3'$. Since E_1 and E_3' have equal classes, we need to consider two possibilities. First when $E_1 \cong E_3'$ and second when $E_1 \ncong E_3'$.
If $E_1 \cong E_3'$, then the image of the map $p \circ \iota$ is either multiple of identity over E_1 or the zero map. If the former case happens it means that the exact sequence on the first row is split contradicting the assumption that E_2 fits in a non-split exact sequence. If the map is given by the zero map, then we can apply the argument in Lemma 7.3 and obtain a contradiction. If $E_1 \ncong E_3'$ then the map $p \circ \iota$ is the zero map and the proof similarly reduces to argument in proof of Lemma 7.3. □

8 $\mathrm{GL}_2(\mathbb{C})$-invariant stratification

We introduce a $\mathrm{GL}_2(\mathbb{C})$-invariant stratification of $\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$ for β satisfying the condition in Assumption 7.1. Note that by Corollary 4.9

$$\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}_p}(\tilde{\tau}) = \left[\frac{\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}_p^{\mathbf{R}}}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right].$$

Moreover, by Corollary 4.8

$$\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}_p^{\mathbf{R}}}(\tilde{\tau}) = \left[\frac{\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$$

where $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$ is defined as in Definition 4.6. In what follows first we give a stratification of the scheme $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$. Then we take the quotient by $\mathrm{GL}(V)$ and our stratification descends to a stratification of $\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}_p^{\mathbf{R}}}(\tilde{\tau})$. Finally we take the quotient of the strata of $\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}_p^{\mathbf{R}}}(\tilde{\tau})$ by the action of $\mathrm{GL}_2(\mathbb{C})$ and we obtain a $\mathrm{GL}_2(\mathbb{C})$-invariant stratification of $\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}_p}(\tilde{\tau})$. The figure below explains our strategy schematically:


$\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$
Stratification
π_1
π_1
$\left[\frac{\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})}{\mathrm{GL}(V)}\right] \cong \mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}^{\mathbf{R}}_p}(\tilde{\tau})$
Induced Stratification
π_2
π_2
$\mathfrak{M}^{(\beta,2)}_{\mathrm{ss},\mathcal{B}_\mathrm{p}}(\tilde{\tau}) \cong \left[\frac{\mathfrak{M}^{(\beta,2)}_{ss,\mathcal{B}^{\mathbf{R}}_p}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right]$
Induced Stratification

Definition 8.1. Define $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$ to be given as the corresponding parametrizing scheme of strictly semistable objects in $\mathcal{B}_p$ of type $(\beta,2)$ (for β as in Assumption 7.1) which are obtained as an extension of two $\tilde{\tau}$-stable objects of rank 1. In other words an object $E_2 \in \mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$ with class $(\beta,2)$ fits in an exact sequence

$$0 \to E_1 \to E_2 \to E_3 \to 0, \tag{8.1}$$

where E_1 and E_3 are $\tilde{\tau}$-stable objects with classes $(\beta_k,1)$ and $(\beta_l,1)$ respectively for some β_k and β_l such that $\beta_k+\beta_l=\beta$.

Remark 8.2. Note that the existence of exact sequence (6.1) and the discussion in Section 6 shows that all objects in $\mathcal{B}_p$ of type $(\beta,2)$ are strictly $\tilde{\tau}$-semistable. Hence $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau}) \cong \mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$ where $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$ is defined in Definition 4.6.

If the extension in (8.1) is non-split, then its automorphism group is obtained by $\mathrm{Hom}(E_3,E_1)\rtimes\mathbb{G}_m$ and if split, the automorphism group is obtained by $\mathrm{Hom}(E_3,E_1)\rtimes\mathbb{G}_m^2$ [9] (page 33). We need these automorphism groups in order to compute the product $(*)$ of the elements of the Ringe-Hall algebra. These elements are given as stack functions which parametrize objects of a given type (such as $(\beta_k,1)$ or $(\beta_l,1)$). Now assume that the exact sequence in (8.1) is non-split and moreover $E_1\cong E_3$. In this case since a semistable object in $\mathfrak{S}^{(\beta,1)}_{st-ss}(\tilde{\tau})$ is also stable, by the property of β in Assumption 7.1 and Lemma 7.2, $\mathrm{Hom}(E_3,E_1)\cong\mathbb{A}^1$ and the automorphism group of extension (8.1) is obtained by $\mathbb{A}^1\rtimes\mathbb{G}_m$. Moreover if in (8.1) $E_1\ncong E_3$, then by Lemma 7.2, $\mathrm{Hom}(E_3,E_1)=0$ and the automorphism group of extension (8.1) is obtained by $\mathbb{G}_m$.

Following similar argument for case of split extensions, we find that the automorphism group of the split extension is $\mathbb{A}^1\rtimes\mathbb{G}_m^2$ when $E_1\cong E_3$ and $\mathbb{G}_m^2$ when $E_1\ncong E_3$. First we decompose $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$

into two disjoint strata:

$$\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau}) = \mathfrak{S}^{(\beta,2)}_{n-sp}(\tilde{\tau}) \bigsqcup \mathfrak{S}^{(\beta,2)}_{sp}(\tilde{\tau}).$$

Here $\mathfrak{S}^{(\beta,2)}_{n-sp,\mathcal{B}_p}(\tilde{\tau})$ and $\mathfrak{S}^{(\beta,2)}_{sp,\mathcal{B}_p}(\tilde{\tau})$ stand for the strata over which the objects representing the elements in $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$ are given by non-split and split extensions respectively. These strata are disjoint since an element $E_2 \in \mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$ can not be given by both split and non-split extensions. By lemmas 7.3 and 7.4, $\mathfrak{S}^{(\beta,2)}_{n-sp}(\tilde{\tau})$ can further be stratified into a disjoint union over all possible β_k and β_l:

$$\mathfrak{S}^{(\beta,2)}_{n-sp}(\tilde{\tau}) = \bigsqcup_{\beta_k+\beta_l=\beta} \mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau}),$$

where $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau})$ stands for stratum over which E_2 is obtained by a non-split extension of $\tilde{\tau}$-stable objects E_1 by E_3 with fixed classes $(\beta_k, 1)$ and $(\beta_l, 1)$ respectively. Now take one of these strata $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau})$ by fixing β_k and β_l. We claim that one may decompose this stratum further into two disjoint strata.

1. Define $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{1}(\tilde{\tau})$ to be the parametrizing scheme of objects $E_2 \in \mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau})$ such that there exists a non-split exact sequence

   $$0 \to E_1 \to E_2 \to E_3 \to 0,$$

   where $E_1 \in \mathfrak{S}^{(\beta_k,1)}_{s}(\tilde{\tau})$ and $E_3 \in \mathfrak{S}^{(\beta_l,1)}_{s}(\tilde{\tau})$ and $E_1 \ncong E_3$.

2. Define $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau})$ to be the parametrizing scheme of objects $E_2 \in \mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau})$ such that there exists a non-split exact sequence

   $$0 \to E_1 \to E_2 \to E_3 \to 0,$$

   where $E_1, E_3 \in \mathfrak{S}^{(\frac{\beta}{2},1)}_{s}(\tilde{\tau})$ and $E_1 \cong E_3$.

Lemma 8.3. *There exists a stratification of* $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau})$*:*

$$\mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau}) = \mathfrak{S}^{(\beta_k,\beta_l,2)}_{1}(\tilde{\tau}) \bigsqcup \mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau}). \tag{8.2}$$

Proof. : We show that $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{1}(\tilde{\tau})$ and $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau})$ are disjoint. Assume that $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{1}(\tilde{\tau}) \bigcap \mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau}) \neq \varnothing$. Therefore, there exist an object $E_2 \in \mathfrak{S}^{(\beta_k,\beta_l,2)}_{n-sp}(\tilde{\tau})$ which fits in exact sequences:

$$0 \to E_1 \to E_2 \to E_3 \to 0,$$

and

$$0 \to E'_1 \to E_2 \to E'_3 \to 0,$$

such that $E_1 \ncong E_3$ and $E'_1 \cong E'_3$. Hence, one obtains a map $p \circ \iota : E_1 \to E'_3$ via the following diagram:

$$\begin{tikzcd}
0 \arrow[r] & E_1 \arrow[r, "\iota"] & E_2 \arrow[r] \arrow[d, "\cong"] & E_3 \arrow[r] & 0 \\
0 \arrow[r] & E'_1 \arrow[r] & E_2 \arrow[r, "p"'] & E'_3 \arrow[r] & 0
\end{tikzcd} \quad . \tag{8.3}$$

If $E_1 \cong E'_3$, then the image of $p \circ \iota$ is multiple of identity over E_1 or the zero map. For the former case we conclude that the first row splits, hence a contradiction. If $p \circ \iota$ is the zero map, then $p \circ \iota$ factors through the map $\iota' \circ g$ in the following diagram:

$$\begin{tikzcd}
0 \arrow[r] & E_1 \arrow[r, "\iota"] \arrow[d, "g"] & E_2 \arrow[r] \arrow[d, "\cong"] & E_3 \arrow[r] & 0 \\
0 \arrow[r] & E'_1 \arrow[r, "\iota'"] & E_2 \arrow[r, "p"'] & E'_3 \arrow[r] & 0
\end{tikzcd} . \tag{8.4}$$

Since $E_1 \cong E'_3$ and $E'_3 \cong E'_1$ then $E_1 \cong E'_1$ and we conclude one of the following possibilities: either ι and ι' are both given as zero maps which is over-ruled (since ι and ι' are both injections) or the map g is zero which is obviously overruled. Moreover if the map g is an isomorphism then $E_3 \cong E'_3$ and since $E_1 \cong E'_3$ by assumption, then $E_1 \cong E_3$ which is a contradiction. Therefore, we assume $E_1 \ncong E'_3$. In this case $p \circ \iota \in \operatorname{Hom}(E_1, E'_3)$ which is the zero map by Lemma 7.2. Then similarly $p \circ \iota$ factors through the map $\iota' \circ g$ in the diagram (8.4). Since $E'_1 \cong E'_3$ and $E_1 \ncong E'_3$, then $E_1 \ncong E'_1$ and by Lemma 7.2, g is the zero map. By considering the left commutative square in (8.4) we obtain a contradiction since the image of E_1 in E_2 is nonzero. Therefore $\mathfrak{S}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \bigcap \mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}) = \varnothing$. $\square$

So far we have proved

$$\mathfrak{S}_{n-sp}^{(\beta,2)}(\tilde{\tau}) = \bigsqcup_{\beta_k+\beta_l=\beta} \left(\mathfrak{S}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \bigsqcup \mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \right).$$

Now consider $\mathfrak{S}_{sp}^{(\beta,2)}(\tilde{\tau})$. We further decompose this stratum as follows.

1. Define $\mathfrak{S}_{\Delta}^{(\beta,2)}(\tilde{\tau})$ to be the parametrizing scheme of objects $E_2 \in \mathfrak{S}_{sp}^{(\beta,2)}(\tilde{\tau})$ such that
$$E_2 \cong E_1 \oplus E_3$$
where $E_1, E_3 \in \mathfrak{S}_s^{(\frac{\beta}{2},1)}(\tilde{\tau})$ and $E_1 \cong E_3$.
2. Define $\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ to be the parametrizing scheme of objects $E_2 \in \mathfrak{S}_{sp}^{(\beta,2)}(\tilde{\tau})$ such that
$$E_2 \cong E_1 \oplus E_3$$
where $E_1 \in \mathfrak{S}_s^{(\beta_k,1)}(\tilde{\tau})$, $E_3 \in \mathfrak{S}_s^{(\beta_l,1)}(\tilde{\tau})$ and $E_1 \ncong E_3$.

By the same argument as in Lemma 8.3 we see that

$$\mathfrak{S}_{sp}^{(\beta,2)}(\tilde{\tau}) = \mathfrak{S}_{\Delta}^{(\beta,2)}(\tilde{\tau}) \bigsqcup \mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau}).$$

The above arguments enable us to give a stratification of $\mathfrak{S}_{st-ss}^{(\beta,2)}(\tilde{\tau})$:

$$\begin{aligned}
&\mathfrak{S}_{st-ss}^{(\beta,2)}(\tilde{\tau}) = \\
&\bigsqcup_{\beta_k+\beta_l=\beta}^{o} \left(\mathfrak{S}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \bigsqcup \mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \right) \bigsqcup \\
&\bigsqcup_{\beta_k+\beta_l=\beta}^{u-o} \mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \bigsqcup \mathfrak{S}_{\Delta}^{(\beta,2)}(\tilde{\tau})
\end{aligned} \tag{8.5}$$

Here $\overset{o}{\underset{\beta_k+\beta_l=\beta}{\bigsqcup}}$ and $\overset{u-o}{\underset{\beta_k+\beta_l=\beta}{\bigsqcup}}$ stand for ordered and un-ordered disjoint unions respectively. We introduce a new notation. Fix β_k and β_l such that $\beta_k+\beta_l=\beta$. Let $\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ be the parametrizing scheme of objects $E_2\cong E_1\oplus E_3$ such that $(E_1,E_3)\in\mathfrak{S}^{(\beta_k,1)}_s(\tilde{\tau})\times\mathfrak{S}^{(\beta_l,1)}_s(\tilde{\tau})$ and $E_1\cong E_3$. It is obvious from the definition that if $\beta_k\neq\beta_l$ then $\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\varnothing$ and for $\beta_k=\beta_l=\frac{\beta}{2}$, $\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\mathfrak{S}^{(\beta,2)}_{\Delta}(\tilde{\tau})$. Using this new notation, $\mathfrak{S}^{(\beta,2)}_{\Delta}(\tilde{\tau})=\underset{\beta_k+\beta_l=\beta}{\bigsqcup}\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ (ordered or unordered unions in this case are same as one) and by (8.5) we obtain:

$$\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})=\overset{o}{\underset{\beta_k+\beta_l=\beta}{\bigsqcup}}\left(\mathfrak{S}^{(\beta_k,\beta_l,2)}_1(\tilde{\tau})\bigsqcup\mathfrak{S}^{(\beta_k,\beta_l,2)}_2(\tilde{\tau})\right)\bigsqcup$$
$$\overset{u-o}{\underset{\beta_k+\beta_l=\beta}{\bigsqcup}}\left(\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})\bigsqcup\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau})\right)\tag{8.6}$$

Lemma 8.4. *Fix (β_k,β_l) such that $\beta_k+\beta_l=\beta$, then the (k,l)'th summand on the right hand side of Equation* (8.6) *is composed of locally closed strata. Moreover $\mathfrak{S}^{(\beta,2)}_s(\tilde{\tau})$ is locally closed.*

Proof. . The (k,l)'th summand on the right hand side of (8.6) is given by:

$$R=\mathfrak{S}^{(\beta_k,\beta_l,2)}_1(\tilde{\tau})\bigsqcup\mathfrak{S}^{(\beta_k,\beta_l,2)}_2(\tilde{\tau})\bigsqcup\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})\bigsqcup\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau}).\tag{8.7}$$

By construction in Section 4.1 and Remark 8.2 there exists an action of $\mathrm{GL}_2(\mathbb{C})\times\mathrm{GL}(V)$ on $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$. This action induces an action of the corresponding Lie algebra on the tangent space of $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$ given by the map:

$$\mathcal{O}_{\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})}\otimes\mathfrak{Lie}(\mathrm{GL}_2(\mathbb{C})\times\mathrm{GL}(V))\to T_{\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})},\tag{8.8}$$

where $\mathfrak{Lie}(\mathrm{GL}_2(\mathbb{C})\times\mathrm{GL}(V))$ denotes the Lie algebra associated to the group $\mathrm{GL}_2(\mathbb{C})\times\mathrm{GL}(V)$. The dimension of the automorphism group of objects representing the elements of $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$ is given by the dimension of the stabilizer (in $\mathrm{GL}_2(\mathbb{C})\times\mathrm{GL}(V)$) group of these elements, which itself is given by the dimension of the kernel of the map in (8.8). On the other hand, the dimension of the kernel of the map in (8.8) is an upper-semicontinious function. Therefore, we obtain a stratification of $\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})$ which induces a stratification of R into locally closed strata such that over each stratum the dimension of the stabilizer group is constant as we vary over points inside that stratum.
The stabilizer group of points in $\mathfrak{S}^{(\beta_k,\beta_l,2)}_1(\tilde{\tau})$ is given by the automorphism group of their corresponding objects (i.e given by extensions of stable non-isomorphic objects with classes $(\beta_k,1)$ and $(\beta_l,1)$) which by above discussions is given by $\mathbb{G}_m$. Let

$$R_1=\mathfrak{S}^{(\beta_k,\beta_l,2)}_1(\tilde{\tau}).$$

By construction, the stabilizer groups of elements in $\mathfrak{S}^{(\beta_k,\beta_l,2)}_2(\tilde{\tau})$ and $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})$ are given by $\mathbb{A}_1\rtimes\mathbb{G}_m$ and $\mathbb{G}^2_m$ respectively. Let

$$R_2=\mathfrak{S}^{(\beta_k,\beta_l,2)}_2(\tilde{\tau})\bigsqcup\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau}).$$

By construction, the stabilizer group of elements in $\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ is given by $\mathbb{A}_1 \rtimes \mathbb{G}_m^2$. Let

$$R_3 = \mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau}).$$

Hence, the right hand side of (8.7) is written as

$$R = R_1 \bigsqcup R_2 \bigsqcup R_3,$$

where the points over R_1, R_2 and R_3 have 1-dimensional, 2-dimensional and 3-dimensional stabilizer groups respectively. It is true that R_1, R_2 and R_3 are locally closed in $\mathfrak{S}_{ss}^{(\beta,2)}(\tilde{\tau})$. Now consider R_2. Let $\overline{\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})}$ denote the closure of $\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ in $\mathfrak{S}_{ss}^{(\beta,2)}(\tilde{\tau})$. Recall that by definition, the objects parametrized by elements of $\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ are given by split extensions, i.e $E_2 \cong E_1 \oplus E_3$ where $E_1 \ncong E_3$. The automorphism group of these objects is given by $\mathbb{G}_m^2$. Taking the closure of $\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$, we immediately see that the objects parametrized by $\overline{\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})}$ are given by all split extensions of E_1 by E_3, i.e one has:

$$\overline{\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})} \subset \mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \bigcup R_3. \tag{8.9}$$

Now take the closure of $\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ and obtain $\overline{\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})}$. By definition, the objects representing elements of $\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ are given by non-split exact sequences

$$0 \to E_1 \to E_2 \to E_3 \to 0,$$

where $E_1 \cong E_3$. So it is seen that taking the closure, the objects representing the elements in the boundary of $\overline{\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})}$ are given by $E_2 \cong E_1 \oplus E_3$ where $E_1 \cong E_3$, i.e:

$$\overline{\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})} \subset \mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \bigcup R_3 \tag{8.10}$$

Since $R_3 \bigcap \mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau}) = \varnothing$ and $R_3 \bigcap \mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}) = \varnothing$, then it is seen that $\overline{\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})}$ and $\overline{\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})}$ have empty intersections in R_2 but non-empty intersections in R_3 , in other words their boundary is given by a subset of R_3 which itself is locally closed in $\mathfrak{S}_{ss}^{(\beta,2)}(\tilde{\tau})$. Hence $\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ and $\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ are locally closed in $\mathfrak{S}_{ss}^{(\beta,2)}(\tilde{\tau})$. $\square$

Remark 8.5. *Here we discuss a second approach to the proof of the fact that* $\mathfrak{S}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ *and* $\mathfrak{S}_{O-\Delta}^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ *are locally closed in* R_2.

Proof. : First we state a theorem from SGA3:

Theorem 8.6. *(SGA3 Exp, X. Theorem 8.8) Let* $\mathcal{T}$ *be a commutative flat group scheme, separated of finite type over a noetherian scheme* S *, with connected affine fibers. Let* $s \in S$ *and* $\overline{s}$ *a geometric point over* s *and suppose:*

1. *the reduced subscheme* $(\mathcal{T}_{\overline{s}})_{red}$ *of the geometric fiber* $\mathcal{T}_{\overline{s}}$ *is a torus and;*
2. *there exists a generization* t *of* s *(i.e. the closure of* $\{t\}$ *contains* s *) such that* $\mathcal{T}_t$ *is smooth over* $k(t)$*, the residue field of* t*;*

Then there exists an open neighborhood $\mathcal{U}$ of s such that $\mathcal{T}\mid_{\mathcal{U}}$ is a torus over $\mathcal{U}$.

Note that , by construction, R_2 is given as a stratum of $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$ over which the stabilizer groups of associated points are two dimensional. As we vary over R_2 the stabilizer groups of points in R_2 make a group scheme $\mathcal{G}$ over $\mathbb{C}$. According to the above theorem for every point $p \in R_2$ such that $(\mathcal{G}_p)_{red}$ is given by the two dimensional torus $(\mathbb{G}^2_m)$ there exists an open neighborhood $\mathcal{U}$ such that $\mathcal{G}\mid_{\mathcal{U}}$ is given by $\mathbb{G}^2_m$ over $\mathcal{U}$. Let $\mathcal{K}$ denote the union of all such $\mathcal{U}$. It is easily seen that every geometric point $k \in \mathcal{K}$ is given as a two dimensional torus $\mathbb{G}^2_m$ which corresponds to the stabilizer group of a point $p \in \mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})$ and there exists a bijective correspondence between such k and p. Hence according to Theorem 8.6, the locus of points $p \in R_2$ with torus stabilizers, i.e $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})$ is open in R_2. Since R_2 is locally closed in $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$ then $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})$ is locally closed in $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$. On the other hand the complement $(\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau}))^c$ is closed in R_2. Since $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})\bigcap\mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau}) = \varnothing$ then $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau}) \subset (\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau}))^c$ which is closed in R_2, hence $\mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau})$ is locally closed in $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$. □

Now consider the (k,l)'th summand on the right hand side of (8.6). The action of $\mathrm{GL}(V)$ on $\mathfrak{S}^{(\beta,2)}_{ss}(\tilde{\tau})$ induces an action on each stratum. Take the quotient of each stratum by $\mathrm{GL}(V)$:

Definition 8.7. Define

1. $\mathcal{M}^{(\beta_k,\beta_l,2)}_{1}(\tilde{\tau}) = \left[\frac{\mathfrak{S}^{(\beta_k,\beta_l,2)}_{1}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$.
2. $\mathcal{M}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau}) = \left[\frac{\mathfrak{S}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$.
3. $\mathcal{M}^{(\beta_k,\beta_l,2)}_{3}(\tilde{\tau}) = \left[\frac{\mathfrak{S}^{(\beta_k,\beta_l,2)}_{O-\Delta}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$.
4. $\mathcal{M}^{(\beta_k,\beta_l,2)}_{4}(\tilde{\tau}) = \left[\frac{\mathfrak{D}^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$.

Definition 8.8. Define $\mathcal{M}^{(\beta,2)}_{st-ss}(\tilde{\tau}) = \left[\frac{\mathfrak{S}^{(\beta,2)}_{st-ss}(\tilde{\tau})}{\mathrm{GL}(V)}\right]$.

By definitions 8.7 and 8.8 and Equation (8.6) one obtains:

$$\mathcal{M}^{(\beta,2)}_{st-ss}(\tilde{\tau}) = \coprod^{o}_{\beta_k+\beta_l=\beta}\left(\mathcal{M}^{(\beta_k,\beta_l,2)}_{1}(\tilde{\tau})\coprod\mathcal{M}^{(\beta_k,\beta_l,2)}_{2}(\tilde{\tau})\right)\coprod$$
$$\coprod^{u-o}_{\beta_k+\beta_l=\beta}\left(\mathcal{M}^{(\beta_k,\beta_l,2)}_{3}(\tilde{\tau})\coprod\mathcal{M}^{(\beta_k,\beta_l,2)}_{4}(\tilde{\tau})\right) \tag{8.11}$$

Moreover by Corollary 4.8 and Remark 8.2:

$$\mathfrak{M}^{(\beta,2)}_{\mathcal{B}^{\mathbf{R}}_p,ss}(\tilde{\tau}) = \mathcal{M}^{(\beta,2)}_{st-ss}(\tilde{\tau}). \tag{8.12}$$

Now take the quotient of $\mathfrak{M}^{(\beta,2)}_{\mathcal{B}^{\mathbf{R}}_p,ss}(\tilde{\tau})$ by the action of $\mathrm{GL}_2(\mathbb{C})$ and obtain $\mathfrak{M}^{(\beta,2)}_{\mathcal{B}_p,ss}(\tilde{\tau})$. The action of $\mathrm{GL}_2(\mathbb{C})$ restricts to each stratum, hence one may take the quotient of each stratum by $\mathrm{GL}_2(\mathbb{C})$ and define:

Definition 8.9. Define

1. $\mathfrak{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}) = \left[\frac{\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right]$.
2. $\mathfrak{M}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}) = \left[\frac{\mathcal{M}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right]$.
3. $\mathfrak{M}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau}) = \left[\frac{\mathcal{M}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right]$.
4. $\mathfrak{M}_4^{(\beta_k,\beta_l,2)}(\tilde{\tau}) = \left[\frac{\mathcal{M}_4^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right]$.

Now we associate characteristic stack functions to each stacky stratum.

Definition 8.10. Define

1. $\bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ to be the characteristic stack function associated to $\mathfrak{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})$.
2. $\bar{\delta}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ to be the characteristic stack function associated to $\mathfrak{M}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})$.
3. $\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ to be the characteristic stack function associated to $\mathfrak{M}_3^{(\beta_l,2)}(\tilde{\tau}))$.
4. $\bar{\delta}_4^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ to be the characteristic stack function associated to $\mathfrak{M}_4^{(\beta_k,\beta_l,2)}$.

Definition 8.11. Define

$$\mathfrak{M}_{st-ss}^{(\beta,2)}(\tilde{\tau}) = \left[\frac{\mathcal{M}_{st-ss}^{(\beta,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right] \tag{8.13}$$

By by Corollary 4.9, Corollary 4.8 and Equation (8.12) it is easily seen that

$$\mathfrak{M}_{\mathcal{B}_p,ss}^{(\beta,2)}(\tilde{\tau}) = \mathfrak{M}_{st-ss}^{(\beta,2)}(\tilde{\tau}).$$

Definition 8.12. Define $\bar{\delta}_{st-ss}^{(\beta,2)}(\tilde{\tau})$ to be the characteristic stack function of $\mathfrak{M}_{st-ss}^{(\beta,2)}(\tilde{\tau})$. Moreover define $\bar{\delta}_s^{(\beta,2)}(\tilde{\tau})$ to be the characteristic stack function $\mathfrak{M}_s^{(\beta,2)}(\tilde{\tau})$.

Definition 8.13. Define $\bar{\delta}_{ss}^{(\beta,2)}(\tilde{\tau})$ to be the characteristic stack function of $\mathfrak{M}_{\mathcal{B}_p,ss}^{(\beta,2)}(\tilde{\tau})$.

Hence by definitions 8.12 and 8.13 the following identity holds true:

$$\bar{\delta}_{ss}^{(\beta,2)}(\tilde{\tau}) = \bar{\delta}_{st-ss}^{(\beta,2)}(\tilde{\tau}). \tag{8.14}$$

Now by Equation (8.11) and Definition 8.10 obtain:

$$\bar{\delta}_{st-ss}^{(\beta,2)}(\tilde{\tau}) = \sum_{\beta_k+\beta_l=\beta}^{o} \left(\bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}) + \bar{\delta}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})\right) + \sum_{\beta_k+\beta_l=\beta}^{u-o} \left(\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau}) + \bar{\delta}_4^{(\beta_k,\beta_l,2)}(\tilde{\tau})\right), \tag{8.15}$$

where $\sum_{\beta_k+\beta_l=\beta}^{o}$ and $\sum_{\beta_k+\beta_l=\beta}^{u-o}$ denote the ordered and un-ordered sums respectively. Therefore, by

(8.15) and (8.14) the following identity holds true:

$$\bar{\delta}_{ss}^{(\beta,2)}(\tilde{\tau}) = \sum_{\beta_k+\beta_l=\beta}^{o}\Big(\bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})+\bar{\delta}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})\Big)+\sum_{\beta_k+\beta_l=\beta}^{u-o}\Big(\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})+\bar{\delta}_4^{(\beta_k,\beta_l,2)}(\tilde{\tau})\Big). \tag{8.16}$$

However $\bar{\delta}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\varnothing$ and $\bar{\delta}_4^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\varnothing$ for $\beta_k\neq\beta_l$, hence we obtain :

$$\bar{\delta}_{ss}^{(\beta,2)}(\tilde{\tau}) = \sum_{\beta_k+\beta_l=\beta}^{o}\Big(\bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})\Big)+\bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})+\sum_{\beta_k+\beta_l=\beta}^{u-o}\Big(\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})\Big)+\bar{\delta}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}). \tag{8.17}$$

On the other hand by applying Definition 5.5 to $\mathfrak{M}_{ss,\mathcal{B}_p}^{(\beta,2)}(\tilde{\tau})$ we obtain:

$$\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})=\bar{\delta}_{ss}^{(\beta,2)}(\tilde{\tau})-\sum_{\beta_k+\beta_l=\beta}^{o}\frac{1}{2}\bar{\delta}_{ss}^{(\beta_k,1)}(\tilde{\tau})*\bar{\delta}_{ss}^{(\beta_l,1)}(\tilde{\tau}). \tag{8.18}$$

Note that given any class, β_i, there does not exist any strictly semistable object with class $(\beta_i,1)$. Therefore instead of notation $\bar{\delta}_{ss}^{(\beta_k,1)}(\tilde{\tau})$, we use $\bar{\delta}_{s}^{(\beta_k,1)}(\tilde{\tau})$. By equations (8.14), (8.17) and (8.18) we obtain:

$$\begin{aligned}&\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})=\\&\sum_{\beta_k+\beta_l=\beta}^{o}\Big(\bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})\Big)+\bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})+\sum_{\beta_k+\beta_l=\beta}^{u-o}\Big(\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})\Big)+\bar{\delta}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\\&-\sum_{\beta_k+\beta_l=\beta}^{o}\frac{1}{2}\bar{\delta}_{s}^{(\beta_k,1)}(\tilde{\tau})*\bar{\delta}_{s}^{(\beta_l,1)}(\tilde{\tau})\end{aligned} \tag{8.19}$$

It is easily seen that:

$$\sum_{\beta_k+\beta_l=\beta}^{u-o}\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\bar{\delta}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})+\frac{1}{2}\cdot\sum_{\substack{\beta_k+\beta_l=\beta\\ \beta_k\neq\beta_l}}^{o}\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau}).$$

Hence, we rewrite each series on the right hand side of (8.19) as an ordered sum and obtain:

$$\begin{aligned}&\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})=\\&\sum_{\beta_k+\beta_l=\beta}^{o}\Big(\bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})\Big)+\bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})+\bar{\delta}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})+\frac{1}{2}\cdot\sum_{\substack{\beta_k+\beta_l=\beta\\ \beta_k\neq\beta_l}}^{o}\bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})+\bar{\delta}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\\&-\sum_{\beta_k+\beta_l=\beta}^{o}\frac{1}{2}\cdot\bar{\delta}_{s}^{(\beta_k,1)}(\tilde{\tau})*\bar{\delta}_{s}^{(\beta_l,1)}(\tilde{\tau}).\end{aligned} \tag{8.20}$$

Next we compute the ordered product $\bar{\delta}_{s}^{(\beta_k,1)}(\tilde{\tau})*\bar{\delta}_{s}^{(\beta_l,1)}(\tilde{\tau})$ for a fixed choice of β_k and β_l.

9 Computation of $\bar{\delta}_s^{(\beta_k,1)}(\tilde{\tau}) * \bar{\delta}_s^{(\beta_l,1)}(\tilde{\tau})$

In this section we describe the computation of the Ringel hall product of the stack functions $\bar{\delta}_s^{(\beta_k,1)}(\tilde{\tau}) * \bar{\delta}_s^{(\beta_l,1)}(\tilde{\tau})$ for β_k and β_l satisfying the condition in Assumption 7.1. Similar to discussions in Section 5 let $\pi_i : \mathfrak{Exact}_{\mathcal{B}_p} \to \mathfrak{M}_{B_p}(\tilde{\tau})$ for $i = 1, 2, 3$ be the projection map that sends an exact sequence to its first,second and third objects respectively over moduli stack of objects in $\mathcal{B}_p$. We also have the map $\pi_1 \times \pi_3 : \mathfrak{Exact}_{\mathcal{B}_p} \to \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) \times \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau})$. By Joyce's definition in [9]:

$$\delta_{ss}^{(\beta_k,1)}(\tilde{\tau}) * \delta_{ss}^{(\beta_l,1)}(\tilde{\tau}) = \pi_2 * ((\pi_1 \times \pi_3)^*(\delta^{(\beta_k,1)}(\tilde{\tau}) \otimes \delta^{(\beta_l,1)}(\tilde{\tau}))) \tag{9.1}$$

Suppose that $\delta^{(\beta_k,1)} = [\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})/\mathbb{G}_m, \rho_1]$ and $\delta^{(\beta_l,1)} = [\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})/\mathbb{G}_m, \rho_3]$ where $\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})$ and $\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})$ denote some underlying parameter stacks and

$$\rho_1 : \left[\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})/\mathbb{G}_m\right] \to \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}),$$

and

$$\rho_3 : \left[\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})/\mathbb{G}_m\right] \to \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}).$$

Let us denote by $\mathcal{Z}'$ the fibered product

$$(\left[\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})/\mathbb{G}_m\right] \times \left[\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})/\mathbb{G}_m\right]) \times_{\rho_1 \times \rho_3, \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) \times \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}), \pi_1 \times \pi_3} \mathfrak{Exact}_{\mathcal{B}_p}$$

the identity in (9.1) is described by the following diagram:

$$\begin{tikzcd}
\mathcal{Z}' \arrow[r, "\Phi"] \arrow[d] & \mathfrak{Exact}_{\mathcal{B}_p} \arrow[r, "\pi_2"] \arrow[d, "\pi_1 \times \pi_3"] & \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) \\
\left[\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})/\mathbb{G}_m\right] \times \left[\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})/\mathbb{G}_m\right] \arrow[r, "\rho_1 \times \rho_1"] & \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) \times \mathfrak{M}_{\mathcal{B}_p}(\tilde{\tau}) &
\end{tikzcd} \tag{9.2}$$

We compute the product of stack functions in (9.1) by computing it over the $\mathbb{C}$-points of $\bar{\delta}^{(\beta_k,1)}(\tilde{\tau})$ and $\bar{\delta}^{(\beta_l,1)}(\tilde{\tau})$ (these are induced from $\mathbb{C}$-points of $\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})$ and $\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})$) and then integrating over all points in $\mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})$.

Consider the stack function

$$\delta_1 = \left(\left[\frac{\operatorname{Spec}(\mathbb{C})}{\mathbb{G}_m}\right], \rho_1 \circ \iota_1\right),$$

with

$$\iota_1 : \left[\frac{\operatorname{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \to \left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})}{\mathbb{G}_m}\right].$$

Moreover let

$$\delta_3 = \left(\left[\frac{\operatorname{Spec}(\mathbb{C})}{\mathbb{G}_m}\right], \rho_3 \circ \iota_3\right),$$

with

$$\iota_3 : \left[\frac{\operatorname{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] \to \left[\frac{\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}{\mathbb{G}_m}\right].$$

Note that δ_1 and δ_3 are the stack functions associated to $\mathbb{C}$-points of $\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})$ and $\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})$ respectively. Let $E_1 \in \mathfrak{M}^{(\beta_k,1)}_{\mathcal{B}_p}(\tilde{\tau})$ and $E_3 \in \mathfrak{M}^{(\beta_l,1)}_{\mathcal{B}_p}(\tilde{\tau})$ and $E_2 \in \mathfrak{M}^{(\beta,2)}_{\mathcal{B}_p}(\tilde{\tau})$. Consider the exact sequence in $\mathfrak{Exact}_{\mathcal{B}_p}$:

$$0 \to E_1 \to E_2 \to E_3 \to 0 \tag{9.3}$$

The automorphism group of the extension 9.3 is given by $\mathrm{Hom}(E_3, E_1) \rtimes \mathbb{G}_m^2$. The element $(g_1, g_2) \in \mathbb{G}_m^2$ acts on $\mathrm{Ext}^1(E_3, E_1)$ by multiplication by $g_2^{-1} g_1$ and the action of $\mathrm{Hom}(E_3, E_1)$ on $\mathrm{Ext}^1(E_3, E_1)$ is trivial. If the extensions in (9.3) are non-split, then the parametrizing scheme of such extensions is obtained by $\mathbb{P}(\mathrm{Ext}^1(E_3, E_1))$ and for split extensions, it is obtained by $\mathrm{Spec}(\mathbb{C})$. In case of nonsplit extensions, the stabilizer group of the action of $\mathbb{G}_m^2$ is given by $\mathbb{G}_m$ and for split extensions, the stabilizer group of the action of $\mathbb{G}_m^2$ is $\mathbb{G}_m^2$ itself, hence:

$$\delta_1 * \delta_3 = \left(\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathrm{Hom}(E_3, E_1) \rtimes \mathbb{G}_m^2} \right], \mu_1 \right) + \left(\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3, E_1))}{\mathrm{Hom}(E_3, E_1) \rtimes \mathbb{G}_m} \right], \mu_3 \right). \tag{9.4}$$

Definition 9.1. Let $\mathfrak{R}$ be a $\mathbb{C}$-stack. Consider $\mathcal{B}G$ given as a quotient stack $\left[\frac{\mathrm{Spec}(\mathbb{C})}{G}\right]$. Define motivic integration over $\mathfrak{R}$ as an identity in the motivic ring of stack functions:

$$\int_{\mathfrak{R}} \left[\frac{\mathrm{Spec}(\mathbb{C})}{G} \right] d\mu_m := \left[\frac{\mathfrak{R}}{G} \right]. \tag{9.5}$$

Moreover assume that $\mathbf{P} \to \mathfrak{R}$ is a vector bundle over $\mathfrak{R}$. Then define:

$$\int_{\mathfrak{R}} \left[\frac{\mathbf{P}}{G} \right] d\mu_m := \int_{\mathfrak{R}} \chi(\mathbf{P}) \cdot \left[\frac{\mathrm{Spec}(\mathbb{C})}{G} \right] d\mu_m := \chi(\mathbf{P}) \cdot \left[\frac{\mathfrak{R}}{G} \right], \tag{9.6}$$

where $\chi(\mathbf{P})$ denotes the topological Euler characteristic of $\mathbf{P}$. Here The measure μ_m is the map sending constructible sets on $\mathfrak{R}$ to the their corresponding elements in the Grothendieck group of stacks.

Now integrate Equation (9.4) over $\mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})$:

$$\begin{aligned} &\bar{\delta}_s^{(\beta_k,1)}(\tilde{\tau}) * \bar{\delta}_s^{(\beta_l,1)}(\tilde{\tau}) = \int_{(E_1,E_3)\in \mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})} \delta_1 * \delta_3 = \\ &\int_{(E_1,E_3)\in \mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})} \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathrm{Hom}(E_3, E_1) \rtimes \mathbb{G}_m^2} \right] d\mu_m \\ &+ \int_{(E_1,E_3)\in \mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3, E_1))}{\mathrm{Hom}(E_3, E_1) \rtimes \mathbb{G}_m} \right] d\mu_m. \end{aligned} \tag{9.7}$$

By the result of Lemma 7.2, if $\iota_1 = \iota_3$, i.e if $E_1 \cong E_3$, then $\mathrm{Hom}(E_3, E_1) \cong \mathbb{A}^1$ and if $\iota_1 \neq \iota_3$, then

$\mathrm{Hom}(E_3,E_1)=\mathrm{Spec}(\mathbb{C})$, hence:

$$\begin{aligned}&\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathrm{Hom}(E_3,E_1)\rtimes\mathbb{G}_m^2}\right]d\mu_m=\\ &\int_{\Delta}\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{A}^1\rtimes\mathbb{G}_m^2}\right]d\mu_m+\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m^2}\right]d\mu_m\\ &=\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{A}^1\rtimes\mathbb{G}_m^2}\right]+\left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right]\end{aligned}\tag{9.8}$$

and similarly:

$$\begin{aligned}&\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathrm{Hom}(E_3,E_1)\rtimes\mathbb{G}_m}\right]d\mu_m\\ &=\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{A}^1\rtimes\mathbb{G}_m}\right]d\mu_m+\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right]d\mu_m.\end{aligned}\tag{9.9}$$

From Equations (9.7), (9.8) and (9.9) we obtain:

$$\begin{aligned}&\bar{\delta}_s^{(\beta_k,1)}(\tilde{\tau})*\bar{\delta}_s^{(\beta_l,1)}(\tilde{\tau})=\\ &\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{A}^1\rtimes\mathbb{G}_m^2}\right]+\left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right]+\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{A}^1\rtimes\mathbb{G}_m}\right]d\mu_m\\ &+\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right]d\mu_m.\end{aligned}\tag{9.10}$$

Adding the contribution of the terms obtained on the right hand side of Equation (9.10) for all possible choices of β_k and β_l one obtains:

$$\begin{aligned}&\sum_{\beta_k+\beta_l=\beta}^{o}\bar{\delta}_s^{(\beta_k,1)}(\tilde{\tau})*\bar{\delta}_s^{(\beta_l,1)}(\tilde{\tau})=\sum_{\beta_k+\beta_l=\beta}^{o}\left(\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right]d\mu_m\right)\\ &+\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{A}^1\rtimes\mathbb{G}_m}\right]d\mu_m+\sum_{\beta_k+\beta_l=\beta}^{o}\left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right]+\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{A}^1\rtimes\mathbb{G}_m^2}\right]\end{aligned}\tag{9.11}$$

It is easily seen that

$$\begin{aligned}&\sum_{\beta_k+\beta_l=\beta}^{o}\left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right]=\\ &\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right]+\sum_{\substack{\beta_k+\beta_l=\beta\\ \beta_k\neq\beta_l}}^{o}\left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right]\end{aligned}\tag{9.12}$$

Hence

$$\begin{aligned}
&\sum_{\beta_k+\beta_l=\beta}^{o} \bar{\delta}_s^{(\beta_k,1)}(\tilde{\tau}) * \bar{\delta}_s^{(\beta_l,1)}(\tilde{\tau}) = \\
&\sum_{\beta_k+\beta_l=\beta}^{o} \left(\int_{(E_1,E_3)\in \mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right] d\mu_m \right) \\
&+ \int_{E_1\in \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{A}^1 \rtimes \mathbb{G}_m}\right] d\mu_m + \left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right] \\
&+ \sum_{\substack{\beta_k+\beta_l=\beta \\ \beta_k\neq\beta_l}}^{o} \left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right] + \left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{A}^1\rtimes \mathbb{G}_m^2}\right].
\end{aligned} \tag{9.13}$$

By Equations (9.13), (8.20), we obtain:

$$\begin{aligned}
\bar{\epsilon}^{(\beta,2)}(\tilde{\tau}) = &\sum_{\beta_k+\beta_l=\beta}^{o} \left(\bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}) - \frac{1}{2}\int_{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right] d\mu_m \right) \\
&+ \left(\bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) - \frac{1}{2}\int_{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{A}^1\rtimes \mathbb{G}_m}\right] d\mu_m \right) \\
&+ \left(\bar{\delta}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) - \frac{1}{2}\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right] \right) \\
&+ \sum_{\substack{\beta_k+\beta_l=\beta \\ \beta_k\neq\beta_l}}^{o} \left(\frac{1}{2}\cdot \bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau}) - \frac{1}{2}\left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right] \right) + \left(\bar{\delta}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) - \frac{1}{2}\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{A}^1\rtimes \mathbb{G}_m^2}\right] \right).
\end{aligned} \tag{9.14}$$

We consider a new notation for the right hand side of Equation (9.14). Let

$$\begin{aligned}
\epsilon_1^{(k,l)}(\tilde{\tau}) &= \bar{\delta}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}) - \frac{1}{2}\int_{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times \mathfrak{S}^{(\beta_l,1)}\backslash\Delta} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right] d\mu_m \\
\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) &= \bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) - \frac{1}{2}\int_{\mathcal{M}^{(\frac{\beta}{2},1)}} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{A}^1\rtimes \mathbb{G}_m}\right] d\mu_m \\
\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) &= \bar{\delta}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) - \frac{1}{2}\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right] \\
\epsilon_3^{(k,l)}(\tilde{\tau}) &= \left(\frac{1}{2}\cdot \bar{\delta}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau}) - \frac{1}{2}\left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right] \right) |_{\beta_k\neq\beta_l} \\
\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) &= \bar{\delta}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) - \frac{1}{2}\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{A}^1\rtimes \mathbb{G}_m^2}\right].
\end{aligned} \tag{9.15}$$

Using this notation, Equation (9.14) is rewritten as:

$$\bar{\epsilon}^{(\beta,2)}(\tilde{\tau})=\sum_{\beta_k+\beta_l=\beta}^{o}\epsilon_1^{(k,l)}(\tilde{\tau})+\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})+\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})+\sum_{\substack{\beta_k+\beta_l=\beta\\ \beta_k\neq\beta_l}}^{o}\epsilon_3^{(k,l)}(\tilde{\tau})+\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) \tag{9.16}$$

Next we compute the image of Lie algebra morphism, $\tilde{\Psi}^{\mathcal{B}_p}$, applied to $\epsilon_1^{(k,l)}(\tilde{\tau}),\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}),\cdots,\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})$. In order to apply $\tilde{\Psi}^{\mathcal{B}_p}$, one needs to prove that these functions, as elements of the Ringel-Hall algebra, are supported over the virtual indecomposables. Note that as we showed earlier:

$$\bar{\delta}_i^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\left[\frac{\mathcal{M}_i^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right].$$

Joyce in [5] (Section 6.2) has shown that given a stack function $\left[\left(\left[\frac{\mathcal{U}}{\mathrm{GL}_2(\mathbb{C})}\right],\nu\right)\right]$ one has the following identity of stack functions:

$$\begin{aligned}&\left[\left(\left[\frac{\mathcal{U}}{\mathrm{GL}_2(\mathbb{C})}\right],a\right)\right]=\\&F(\mathrm{GL}_2(\mathbb{C}),\mathbb{G}_m^2,\mathbb{G}_m^2)\left[\left(\left[\frac{\mathcal{U}}{\mathbb{G}_m^2}\right],\mu\circ i_1\right)\right]+F(\mathrm{GL}_2(\mathbb{C}),\mathbb{G}_m^2,\mathbb{G}_m)\left[\left(\left[\frac{\mathcal{U}}{\mathbb{G}_m}\right],\mu\circ i_2\right)\right],\end{aligned} \tag{9.17}$$

where

$$\begin{aligned}F(\mathrm{GL}_2(\mathbb{C}),\mathbb{G}_m^2,\mathbb{G}_m^2)&=\frac{1}{2}\\F(\mathrm{GL}_2(\mathbb{C}),\mathbb{G}_m^2,\mathbb{G}_m)&=-\frac{3}{4},\end{aligned} \tag{9.18}$$

and $\mu\circ i_1$ and $\mu\circ i_2$ are the obvious embeddings. Now apply the result of Joyce [5] (Section 6.2) and obtain a decomposition of $\bar{\delta}_i^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ of the following form:

$$\begin{aligned}&\bar{\delta}_i^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\left[\frac{\mathcal{M}_i^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right]=\\&\frac{1}{2}\left[\left(\left[\frac{\mathcal{M}_i^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right],\mu_i'\circ i_1\right)\right]-\frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_i^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathbb{G}_m}\right],\mu_i'\circ i_2\right)\right],\end{aligned} \tag{9.19}$$

where $\mu_i'\circ i_1$ and $\mu_i'\circ i_2$ are the obvious embeddings.

9.1 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_1^{(k,l)}(\tilde{\tau}))$

By definition, for any fixed choice of k,l, the objects $E_2\in\mathfrak{S}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ fit into non-split exact sequences

$$0\to E_1\to E_2\to E_3\to 0 \tag{9.20}$$

where $E_1 \ncong E_3$. In this case, the automorphism group of the extension (9.20) is given by $\mathbb{G}_m$. Since over $\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})$, there exists an action of $\mathrm{GL}_2(\mathbb{C})$ and the stabilizer group of each point in $\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ is $\mathbb{G}_m$, therefore the $\mathrm{GL}_2(\mathbb{C})$ action reduces to a free action of $\mathrm{PGl}_2(\mathbb{C})$ on $\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})$. Hence it is easy to see that there exists a map

$$\pi_1 : \mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})/\,\mathrm{PGl}_2(\mathbb{C}) \to \mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta$$

which sends E_2 to (E_1, E_3).


$\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})/\,\mathrm{PGL}_2(\mathbb{C})$
π_1
$\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})$
$\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})$
Δ

Consider points $(E_1, E_3) \in \mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta$. Consider the fiber $\pi_1 \mid_{(E_1,E_3)}$ which is given by the set of points $E_2 \in \mathfrak{S}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ which fit in the exact sequence (9.20). Dividing by the automorphism group of extension (9.20), $\mathbb{G}_m$, would provide a bijective correspondence between the set of isomorphism classes of such E_2 (which still undergo a free action of $\mathrm{PGL}_2(\mathbb{C})$) and the set of (E_1, E_3). Therefore there exists a bijective map between the closed points of the fiber of π_1 over $(E_1, E_3) \in \mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta$ and the closed points of $\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})/\,\mathrm{PGL}_2(\mathbb{C})$ which fit into the exact sequence (9.20), i.e the closed points of $\mathbb{P}(\mathrm{Ext}^1(E_3, E_1))$ over $\mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta$. Now rewrite $\bar{\delta}_1^{(\beta_k,\beta_l,2)}$ as:

$$\bar{\delta}_1^{(\beta_k,\beta_l,2)} = \left[\frac{\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right] = \left[\frac{\mathfrak{S}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})/\,\mathrm{PGL}_2(\mathbb{C})}{\mathbb{G}_m}\right] =$$

$$\int_{\mathcal{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau})/\,\mathrm{PGL}_2(\mathbb{C})} \left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right] d\mu_m = \int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta} \left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right] d\mu_m, \tag{9.21}$$

by Equations (9.21) and (9.14) the equation for $\epsilon_1^{(k,l)}(\tilde{\tau})$ is obtained as:

$$\begin{aligned}
\epsilon_1^{(k,l)}(\tilde{\tau}) &= \int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right]d\mu_m \\
&-\frac{1}{2}\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right]d\mu_m \\
&=\frac{1}{2}\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right]d\mu_m.
\end{aligned} \tag{9.22}$$

Now apply the Lie algebra morphism $\tilde{\Psi}^{\mathcal{B}_p}$ to $\epsilon_1^{(k,l)}(\tilde{\tau})$ and obtain:

$$\begin{aligned}
&\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_1^{(k,l)}(\tilde{\tau})) = \\
&\frac{1}{2}\cdot\chi^{na}\Big(\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))}{\mathbb{G}_m}\right]d\mu_m,\mu^*\nu_{\mathfrak{M}_1^{(\beta,2)}}\Big)\cdot\tilde{\lambda}^{(\beta,2)} \\
&\frac{1}{2}\cdot\Big(\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\chi(\mathbb{P}(\mathrm{Ext}^1(E_3,E_1)))\cdot\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]d\chi\Big)\cdot\tilde{\lambda}^{(\beta,2)} \\
&=(-1)^1\cdot(-1)^{\dim(\mathfrak{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}))-1}\cdot\frac{1}{2}\cdot\Big(\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\chi(\mathbb{P}(\mathrm{Ext}^1(E_3,E_1)))d\chi\Big)\cdot\tilde{\lambda}^{(\beta,2)}
\end{aligned} \tag{9.23}$$

The factor of $(-1)^{\dim(\mathfrak{M}^{(\beta,2)}_{st-ss,\mathcal{B}_p}-1)}$ is due to the fact that a stacky point given as $\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]$ has relative dimension $\dim(\mathfrak{M}_1^{(\beta_k,\beta_l,2)}(\tilde{\tau}))-1$ with respect to the ambient stack. Moreover, the factor of $(-1)^1$ is due the fact that Behrend's function over $\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]$ detects a singular point of order 1.

9.2 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))$

By Equation (9.19)

$$\begin{aligned}
&\bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) = \left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right] = \\
&\frac{1}{2}\left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right],\mu_2'\circ i_1\right)\right]-\frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],\mu_2'\circ i_2\right)\right].
\end{aligned} \tag{9.24}$$

The stabilizer group of points in $\mathfrak{M}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau})=\mathfrak{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ is given by the automorphism group of extensions associated to those points. Since a point $p\in\mathfrak{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})(\mathbb{C})$ is represented by an object E_2 which fits into a non-split exact sequence:

$$0\to E_1\to E_2\to E_1\to 0, \tag{9.25}$$

then the automorphism group of extension (9.25) is given by $\mathbb{A}^1 \rtimes \mathbb{G}_m$. The stabilizer group (at point p) of the action of $\mathrm{GL}_2(\mathbb{C})$ on $\mathcal{M}_2^{(\beta,2)}(\tilde{\tau})$ is obtained by $\mathbb{A}^1 \rtimes \mathbb{G}_m$. Since $E_1 \cong E_3$ then for diagonal matrices (given by $\mathbb{G}_m^2 = T^{\mathrm{GL}_2(\mathbb{C})}$) we have that

$$\mathbb{G}_m^2 \cap \mathrm{Stab}_p(\mathrm{GL}_2(\mathbb{C})) = \mathbb{G}_m \subset \mathbb{G}_m^2,$$

hence the action of $\mathbb{G}_m^2$ descends to a free action of $\mathbb{G}_m^2/\mathbb{G}_m \cong \mathbb{G}_m$. Hence:

$$\left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right], \mu_2' \circ i_1\right)\right] = \left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})/(\mathbb{G}_m^2/\mathbb{G}_m)}{\mathbb{G}_m}\right], \mu_2' \circ i_1\right)\right]. \tag{9.26}$$

Now rewrite $\bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$:

$$\begin{aligned}&\bar{\delta}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}) = \\ &\frac{1}{2}\left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})/(\mathbb{G}_m^2/\mathbb{G}_m)}{\mathbb{G}_m}\right], \mu_2' \circ i_1\right)\right] - \frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right], \mu_2' \circ i_2\right)\right].\end{aligned} \tag{9.27}$$

By equations (9.27) and (9.14):

$$\begin{aligned}&\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) = \\ &\frac{1}{2}\left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})/(\mathbb{G}_m^2/\mathbb{G}_m)}{\mathbb{G}_m}\right], \mu_2' \circ i_1\right)\right] - \frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right], \mu_2' \circ i_2\right)\right] \\ &- \frac{1}{2}\int_{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{G}_m}\right].\end{aligned} \tag{9.28}$$


$\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$
ρ
$\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$
$\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$
Δ

Note that there exists a map from $\rho : \mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)} \to \Delta \cong \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$ which sends an object E_2 fitting in the exact sequence (9.25) to E_1. Moreover, given $E_1 \in \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$, the fiber of ρ over E_1 is the set of points $p \in \mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ represented by exact sequence (9.25). Now for any point $E_1 \in \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$, there exists a map $\pi_{Ext} : \rho^{-1}(\Delta) \to \mathrm{Ext}^1(E_1, E_1)$ which projects the points $p \in \rho^{-1}(E_1)$ to their extension classes. Now consider an element of the extension class $\alpha \in \mathrm{Ext}^1(E_1, E_1)\backslash 0$. The pre-image, $\pi^{-1}_{Ext}(\alpha)$, consists of points $p \in \rho^{-1}(E_1)$ which fall into the class, α. There exists a surjective morphism:

$$\mathrm{GL}_2(\mathbb{C}) \twoheadrightarrow \pi^{-1}_{Ext}(\alpha)$$

which is induced by the $\mathrm{GL}_2(\mathbb{C})$-action on $\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$. The stabilizer group of the action of $\mathrm{GL}_2(\mathbb{C})$ at point $p \in \mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ represented by E_2 sitting inside the exact sequence 9.25 is given by $\mathbb{A}^1 \rtimes \mathbb{G}_m$ hence $\mathrm{Stab}(\mathrm{GL}_2(\mathbb{C}))_p$ for $p \in \rho^{-1}(E_1)$ is given by $\mathbb{A}^1 \rtimes \mathbb{G}_m$, then the fibers of the map $\mathrm{GL}_2(\mathbb{C}) \twoheadrightarrow \pi^{-1}_{Ext}(\alpha)$ are given by $\mathbb{A}^1 \rtimes \mathbb{G}_m$. Now it is easy to relate the virtual poincare polynomial of $\pi^{-1}_{Ext}(\alpha)$ to the virtual Poincaré polynomial of $\mathrm{GL}_2(\mathbb{C})$.
In general given two algebraic spaces X and Y and a fibration $X \to Y$ with fibers Z one has:

$$\mathrm{P}_t(X) = \mathrm{P}_t(Y) \cdot \mathrm{P}_t(Z).$$

Hence we obtain

$$\mathrm{P}_t(\mathrm{GL}_2(\mathbb{C})) = \mathrm{P}_t(\mathbb{A}^1 \rtimes \mathbb{G}_m) \cdot \mathrm{P}_t(\pi^{-1}_{Ext}(\alpha)). \tag{9.29}$$

On the other hand, for each α, the free action of $\mathbb{G}_m^2/\mathbb{G}_m \cong \mathbb{G}_m$ on $\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ induces a free $\mathbb{G}_m$-action on $\pi^{-1}_{Ext}(\alpha)$. Passing to the quotients via this action, we obtain a map

$$\pi^{-1}_{Ext}(\alpha) \to \pi^{-1}_{Ext}(\alpha)/(\mathbb{G}_m^2/\mathbb{G}_m),$$

whose fibers are given by $\mathbb{G}_m^2/\mathbb{G}_m \cong \mathbb{G}_m$. Hence we obtain the following relation for the virtual Poincaré polynomials:

$$\mathrm{P}_t(\pi^{-1}_{Ext}(\alpha)) = \mathrm{P}_t(\mathbb{G}_m^2/\mathbb{G}_m) \cdot \mathrm{P}_t(\pi^{-1}_{Ext}(\alpha)/(\mathbb{G}_m^2/\mathbb{G}_m)). \tag{9.30}$$

By equations (9.29) and (9.30) and according to calculations in [10] (page 4) we obtain:

$$\begin{aligned} \mathrm{P}_t(\pi^{-1}_{Ext}(\alpha)/(\mathbb{G}_m^2/\mathbb{G}_m)) &= \frac{\mathrm{P}_t(\mathrm{GL}_2(\mathbb{C}))}{\mathrm{P}_t(\mathbb{A}^1 \rtimes \mathbb{G}_m) \cdot \mathrm{P}_t(\mathbb{G}_m^2/\mathbb{G}_m)} \\ &= \frac{(t^4-1)\cdot(t^2-1)\cdot t^2}{t^2\cdot(t^2-1)\cdot(t^2-1)} = t^2+1. \end{aligned} \tag{9.31}$$

The computation in Equation (9.31) means that for each $E_1 \in \mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$ by passing to the quotients via the action of $\mathbb{G}_m^2/\mathbb{G}_m$, the map π_{Ext} induces a map

$$\pi_{(\mathbb{G}_m^2/\mathbb{G}_m)}\mid_{E_1} : \rho^{-1}(E_1)/(\mathbb{G}_m^2/\mathbb{G}_m) \to \mathbb{P}(\mathrm{Ext}^1(E_1, E_1))$$

whose fibers have virtual Poincaré polynomial as obtained in Equation (9.31). Moreover, the Euler characteristic of the fibers is computed by evaluating their virtual Poincaré polynomial at $t = 1$, hence for every E_1 the fiber of the map $\pi_{(\mathbb{G}_m^2/\mathbb{G}_m)}|_{E_1}$ has Euler characteristic equal to 2.

Now back to Equation (9.28). Since $\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})$ is supported over virtual indecomposables, we can apply the Lie algebra homomorphism, $\tilde{\Psi}^{\mathcal{B}_p}$:

$$\begin{aligned}
\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})) &= \frac{1}{2}\cdot\chi^{na}\Bigg(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})/(\mathbb{G}_m^2/\mathbb{G}_m)}{\mathbb{G}_m}\right],(\mu_2'\circ i_1)^*\nu_{\mathfrak{M}_2^{(\beta,2)}}\Bigg)\cdot\tilde{\lambda}^{(\beta,2)}\\
&-\frac{3}{4}\cdot\chi^{na}\Bigg(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],(\mu_2'\circ i_2)^*\nu_{\mathfrak{M}_2^{(\beta,2)}}\Bigg)\cdot\tilde{\lambda}^{(\beta,2)}\\
&-\frac{1}{2}\cdot\chi^{na}\Bigg(\int_{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\left[\frac{\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))}{\mathbb{G}_m}\right]d\mu_m,\mu^*\nu_{\mathfrak{M}_2^{(\beta,2)}}\Bigg)\cdot\tilde{\lambda}^{(\beta,2)}.
\end{aligned} \tag{9.32}$$

Now we compute each term on the right hand side of Equation (9.32) separately.

$$\begin{aligned}
&\frac{1}{2}\cdot\chi^{na}\Bigg(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})/(\mathbb{G}_m^2/\mathbb{G}_m)}{\mathbb{G}_m}\right],(\mu_2'\circ i_1)^*\nu_{\mathfrak{M}_2^{(\beta,2)}}\Bigg)\cdot\tilde{\lambda}^{(\beta,2)} =\\
&\Bigg((-1)^{\dim(\mathfrak{M}^{(\beta,2)}_{st-ss,\mathcal{B}_p})-1}\cdot\frac{1}{2}\cdot\int_{E_2\in\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})/(\mathbb{G}_m^2/\mathbb{G}_m)}1\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]d\chi\Bigg)\cdot\tilde{\lambda}^{(\beta,2)} =\\
&\Bigg((-1)^1\cdot(-1)^{\dim(\mathfrak{M}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}))-1}\cdot\frac{1}{2}\cdot 2\cdot\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\chi\Big(\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))\Big)d\chi\Bigg)\cdot\tilde{\lambda}^{(\beta,2)},
\end{aligned} \tag{9.33}$$

By property (2) of the stack functions in Definition 5.2 we have the identity:

$$\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right] = \chi(\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}))\cdot\left[\frac{\mathrm{Spec}(\mathbb{C})}{\mathbb{G}_m}\right]$$

The action of $\mathbb{G}_m\cong\mathbb{G}_m^2/\mathbb{G}_m$ is free on $\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ and since the $\mathbb{G}_m$-fixed locus via this action is empty we have that

$$\chi(\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}))=0,$$

hence the second term in Equation (9.32) vanishes:

$$-\frac{3}{4}\cdot\chi^{na}\Bigg(\left[\frac{\mathcal{M}_2^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],(\mu_2'\circ i_2)^*\nu_{\mathfrak{M}_2^{(\beta,2)}}\Bigg)\cdot\tilde{\lambda}^{(\beta,2)}=0. \tag{9.34}$$

By equations (9.33) and (9.32) we obtain:

$$\begin{aligned}\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})) =& -\left(\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\chi\Big(\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))\Big)d\chi\right)\cdot\tilde{\lambda}^{(\beta,2)}\\ &-\frac{1}{2}\cdot\left(\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\chi\Big(\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))\Big)d\chi\right)\cdot\tilde{\lambda}^{(\beta,2)}\\ &=-\frac{3}{2}\cdot(-1)^{\dim(\mathfrak{M}_2^{(\beta_k,\beta_l,2)}(\tilde{\tau}))-1}\cdot\left(\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\chi\Big(\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))\Big)d\chi\right)\cdot\tilde{\lambda}^{(\beta,2)}.\end{aligned} \tag{9.35}$$

9.3 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))$

By equations (9.14) and (9.19), re-write $\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})$ as follows:

$$\begin{aligned}\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) = \frac{1}{2}\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right],\mu_3'\circ i_1\right)\right]-\frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],\mu_3'\circ i_2\right)\right]\\ -\frac{1}{2}\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right].\end{aligned} \tag{9.36}$$

The set of $T^{\mathrm{GL}_2(\mathbb{C})}/\mathbb{G}_m$-fixed points is given by the image of the map $f:\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta\to\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ such that for every $E_1\ncong E_3$, sends (E_1,E_3) to $E_1\oplus E_3$. Rewrite

$$\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right]=\left[\frac{\mathrm{Im}(f)}{\mathbb{G}_m^2}\right]+\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\backslash\mathrm{Im}(f)}{\mathbb{G}_m^2}\right], \tag{9.37}$$

by injectivity of the map f:

$$\left[\frac{\mathrm{Im}(f)}{\mathbb{G}_m^2}\right]=\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right] \tag{9.38}$$

hence

$$\begin{aligned}
\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) &= \frac{1}{2}\left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right],\mu_3'\circ i_1\right)\right]+\frac{1}{2}\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\backslash\operatorname{Im}(f)}{\mathbb{G}_m^2}\right],\mu_3'\circ i_1\right)\right]\\
&-\frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],\mu_3'\circ i_2\right)\right]-\frac{1}{2}\left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta}{\mathbb{G}_m^2}\right],\mu'\right)\right]\\
&=\frac{1}{2}\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\backslash\operatorname{Im}(f)}{\mathbb{G}_m^2}\right],\mu_3'\circ i_1\right)\right]-\frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],\mu_3'\circ i_2\right)\right]
\end{aligned} \tag{9.39}$$

There exists a free action of $\mathbb{G}_m^2$ on $\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$. Moreover, there exists a trivial action of $\mathbb{G}_m\subset\mathbb{G}_m^2$ on $\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ where $\mathbb{G}_m$ is given by the subgroup of diagonal matrices as in (**??**). Hence the free $\mathbb{G}_m^2$ action on $\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ is reduced to a free action of $\mathbb{G}_m^2/\mathbb{G}_m\cong\mathbb{G}_m$. Therefore:

$$\begin{aligned}
&\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\backslash\operatorname{Im}(f)}{\mathbb{G}_m^2}\right],\mu_3'\circ i_1\right)\right]=\\
&\left[\left(\left[\frac{\left(\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\backslash\operatorname{Im}(f)\right)/(\mathbb{G}_m^2/\mathbb{G}_m)}{\mathbb{G}_m}\right],\mu_3'\circ i_1\right)\right].
\end{aligned} \tag{9.40}$$

Denote

$$\mathcal{A}:=\left(\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\backslash\operatorname{Im}(f)\right)/(\mathbb{G}_m^2/\mathbb{G}_m).$$

The scheme $\mathcal{A}$ is the analogue of $\tilde{Q}_2^{(2,2m)}$ in [16] (Lemma 5.6) when the sheaves F composing the objects $E:=(F,V,\phi)$ have zero dimensional support with length $2m$. In that situation (roughly speaking), the stack $\mathcal{M}_3^{(\beta,2)}(\tilde{\tau})$ is replaced by a subscheme of a product of Quot schemes.
Rewrite $\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})$ as:

$$\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})=\frac{1}{2}\left[\left(\left[\frac{\mathcal{A}}{\mathbb{G}_m}\right],\mu_3'\circ i_1\right)\right]-\frac{3}{4}\left[\left(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],\mu_3'\circ i_2\right)\right]. \tag{9.41}$$

Since $\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})$ is supported over the virtual indecomposables, one can apply the Lie algebra morphism $\tilde{\Psi}^{\mathcal{B}_p}$:

$$\begin{aligned}
&\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})) = \\
&\frac{1}{2}\cdot\chi^{na}\Bigg(\left[\frac{\mathcal{A}}{\mathbb{G}_m}\right],(\mu_3'\circ i_1)^*\nu_{\mathfrak{M}_3^{(\beta,3)}}\Bigg)\cdot\tilde{\lambda}^{(\beta,2)}-\frac{3}{4}\cdot\chi^{na}\Bigg(\left[\frac{\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right],(\mu_3'\circ i_2)^*\nu_{\mathfrak{M}_3^{(\beta,3)}}\Bigg)\cdot\tilde{\lambda}^{(\beta,2)} \\
&=\Bigg((-1)^1\cdot\frac{1}{2}\cdot\chi(\mathcal{A})-(-1)^1\cdot\frac{3}{4}\cdot\chi(\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau}))\Bigg)\cdot\tilde{\lambda}^{(\beta,2)}.
\end{aligned}\tag{9.42}$$

There exists a map $\pi_{13}:\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\to\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta$ which sends a point $E_2\cong E_1\oplus E_3$ to (E_1,E_3). Fix such $(E_1,E_3)\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\backslash\Delta$. The set of points in $\pi_{13}^{-1}(E_1,E_3)$ are those E_2 which can be written as $E_2\cong E_1\oplus E_3$. Moreover, every point in $\pi_{13}^{-1}(E_1,E_3)\mid_{\mathrm{Im}(f)}$ can be determined by $E_2\cong E_1\oplus E_3$ or $E_2\cong E_3\oplus E_1$, i.e by a permutation and an isomorphism, hence there exists an induced map $\pi_{13}':\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})\to\mathrm{Sym}(\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau}))$. The action of $\mathrm{GL}_2(\mathbb{C})$ on $\mathcal{M}_3^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ is restricted to an action on the fiber of π_{13}' over (E_1,E_3).


$\mathcal{A}$
$\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$
$\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$
Δ

The stabilizer group of the points in $(\pi_{13}')^{-1}(E_1,E_3)$ is given by $\mathbb{G}_m^2$. Hence, one concludes that for every $(E_1,E_3)\in\mathrm{Sym}(\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})\times\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau}))$ there exists a map $\mathrm{GL}_2(\mathbb{C})\to\pi_{13}^{-1}(E_1,E_3)$ whose fiber over each point $p\in(\pi_{13}')^{-1}(E_1,E_3)$ is given by $\mathbb{G}_m^2$. Hence, over $(\pi_{13}')^{-1}(E_1,E_3)\backslash\mathrm{Im}(f)$, the action of $\mathrm{GL}_2(\mathbb{C})$ is restricted to $\mathrm{GL}_2(\mathbb{C})\backslash(\mathbb{G}_m^2\bigcup(\mathbb{G}_m^2)^*)$ where $\mathbb{G}_m^2$ is given by the diagonal matrices of the form:

$$\mathbb{G}_m^2=\Bigg\{\begin{pmatrix} g_1 & 0 \\ 0 & g_2\end{pmatrix}\mid g_1,g_2\in\mathbb{C}^*\Bigg\},$$

and $(\mathbb{G}_m^2)^*$ is given by the anti-diagonal matrices of the form:

$$(\mathbb{G}_m^2)^* = \left\{ \left(\begin{array}{cc} 0 & g_1 \\ g_2 & 0 \end{array} \right) \mid g_1, g_2 \in \mathbb{C}^* \right\}.$$

Therefore, there exists a map $\mathrm{GL}_2(\mathbb{C}) \backslash (\mathbb{G}_m^2 \bigcup (\mathbb{G}_m^2)^*) \to (\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f)$ whose fiber over each point $p \in (\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f)$ is given by $\mathbb{G}_m^2$. Now compute the virtual Poincaré polynomial of $(\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f)$.

$$\begin{aligned} \mathrm{P}_t((\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f)) &= \frac{\mathrm{P}_t(\mathrm{GL}_2(\mathbb{C}) \backslash (\mathbb{G}_m^2 \bigcup (\mathbb{G}_m^2)^*))}{\mathrm{P}_t(\mathbb{G}_m^2)} \\ = \frac{(t^4 - 1) \cdot (t^2 - 1) \cdot t^2 - (t^2 - 1)^2}{(t^2 - 1)^2} &= t^4 + t^2 - 1. \end{aligned} \tag{9.43}$$

The occurrence of the term $(t^2 - 1)^2$ in the numerator is due to the free action of $\mathbb{G}_m^2 \bigcup (\mathbb{G}_m^2)^*$ on $\mathrm{Im}(f)$. Note that $(\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f)$ is a $\mathbb{G}_m$ bundle over $(\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f) / \mathbb{G}_m$, hence:

$$\begin{aligned} \mathrm{P}_t((\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f) / \mathbb{G}_m) &= \frac{\mathrm{P}_t((\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f))}{\mathrm{P}_t(\mathbb{G}_m)} \\ = \frac{t^4 + t^2 - 1}{(t^2 - 1)} &= t^2 + 2. \end{aligned} \tag{9.44}$$

Therefore, for every $(E_1, E_3) \in \mathrm{Sym}(\mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}))$ via evaluating the result in Equation (9.44) at $t = 1$, the fibers $(\pi'_{13})^{-1}(E_1, E_3) \backslash \mathrm{Im}(f)$ have Euler characteristic 3. On the other hand, there exists a bijective map between the set of points in the $\mathbb{G}_m^2 / \mathbb{G}_m$-fixed locus of $\mathcal{M}_3^{(\frac{\beta}{2}, \frac{\beta}{2}, 2)}(\tilde{\tau})$ and the set of points in $\mathrm{Im}(f) \cong \mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}) \backslash \Delta$. Denote by $d_3 = \dim(\mathfrak{M}_3^{(\beta_k, \beta_l, 3)}(\tilde{\tau})) - 1$. Now rewrite Equation (9.45):

$$\begin{aligned} \tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(\frac{1}{2}, \frac{1}{2})}(\tilde{\tau})) &= \left((-1)^1 \cdot \frac{1}{2} \cdot \chi(\mathcal{A}) - (-1)^1 \cdot \frac{3}{4} \cdot \chi(\mathcal{M}_3^{(\frac{\beta}{2}, \frac{\beta}{2}, 2)}(\tilde{\tau})) \right) \cdot (-1)^{d_3} \cdot \tilde{\lambda}^{(\beta, 2)} \\ &= \left(-\frac{1}{2} \cdot \int_{\mathcal{A}} 1 d\chi_{\mathcal{A}} + \frac{3}{4} \int_{\mathcal{M}_3^{(\frac{\beta}{2}, \frac{\beta}{2}, 2)}(\tilde{\tau})} 1 d\chi_{\mathcal{M}_3^{(\frac{\beta}{2}, \frac{\beta}{2}, 2)}(\tilde{\tau})} \right) \cdot (-1)^{d_3} \cdot \tilde{\lambda}^{(\beta, 2)} = \\ &- \frac{1}{2} \cdot 3 \cdot \left(\int_{\mathrm{Sym}(\mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}))} 1 d\chi_{\mathrm{Sym}(\mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\frac{\beta}{2}, 1)}(\tilde{\tau}))} \right) \cdot (-1)^{d_3} \cdot \tilde{\lambda}^{(\beta, 2)} \\ &+ \frac{3}{4} \cdot \left(\int_{\mathcal{M}^{(\beta_k, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l, 1)}(\tilde{\tau}) \backslash \Delta} 1 d\chi_{\mathfrak{S}^{(\beta_k, 1)} \times \mathfrak{S}^{(\beta_l, 1)} \backslash \Delta} \right) \cdot (-1)^{d_3} \cdot \tilde{\lambda}^{(\beta, 2)} = \\ &- \frac{1}{2} \cdot 3 \cdot \frac{1}{2} \left(\int_{\mathcal{M}^{(\beta_k, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l, 1)}(\tilde{\tau}) \backslash \Delta} 1 d\chi_{\mathcal{M}^{(\beta_k, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l, 1)}(\tilde{\tau}) \backslash \Delta} \right) \cdot (-1)^{d_3} \cdot \tilde{\lambda}^{(\beta, 2)} \\ &+ \frac{3}{4} \cdot \left(\int_{\mathcal{M}^{(\beta_k, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l, 1)}(\tilde{\tau}) \backslash \Delta} 1 d\chi_{\mathcal{M}^{(\beta_k, 1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l, 1)}(\tilde{\tau}) \backslash \Delta} \right) \cdot (-1)^{d_3} \cdot \tilde{\lambda}^{(\beta, 2)} = 0. \end{aligned} \tag{9.45}$$

9.4 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(k,l)}(\tilde{\tau}))$

Throughout this subsection by earlier construction $\beta_k \neq \beta_l$ and $\beta_k + \beta_l = \beta$. The action of $\mathrm{GL}_2(\mathbb{C})$ restricts to an action of $\mathbb{G}_m^2$ on $\mathcal{M}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})$ hence we obtain

$$\bar{\delta}_3^{(\beta_k,\beta_l)}(\tilde{\tau}) = \left[\frac{\mathcal{M}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right]. \tag{9.46}$$

On the other hand, it is easily seen that there exists a bijective map $f : \mathcal{M}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau}) \to \mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})$ which takes $E_2 \cong E_1 \oplus E_3$ to (E_1, E_3) where $E_1 \ncong E_3$. Hence, we obtain:

$$\left[\frac{\mathcal{M}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right] = \left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right]. \tag{9.47}$$

Therefore, for every pair (β_k, β_l) by definition of $\epsilon_3^{(k,l)}(\tilde{\tau})$ one obtains:

$$\epsilon_3^{(k,l)}(\tilde{\tau}) = \frac{1}{2} \cdot \left(\left[\frac{\mathcal{M}_3^{(\beta_k,\beta_l,2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right] - \left[\frac{\mathcal{M}^{(\beta_k,1)}(\tilde{\tau}) \times \mathcal{M}^{(\beta_l,1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right] \right) = 0, \tag{9.48}$$

hence:

$$\sum_{\substack{\beta_k+\beta_l=\beta \\ \beta_k \neq \beta_l}}^{o} \tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(k,l)}(\tilde{\tau})) = 0. \tag{9.49}$$

9.5 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))$

By construction, there exists a bijective map between the set of points in $\mathcal{M}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})$ and $\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})$, hence one rewrites $\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})$ directly as follows:

$$\begin{aligned}
\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) = \frac{1}{2} \cdot & \left[\left(\left[\frac{\mathcal{M}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m^2}\right], \mu_4' \circ i_1\right)\right] - \frac{3}{4} \cdot \left[\left(\left[\frac{\mathcal{M}_4^{(\frac{\beta}{2},\frac{\beta}{2},2)}(\tilde{\tau})}{\mathbb{G}_m}\right], \mu_4' \circ i_2\right)\right] \\
& - \frac{1}{2} \cdot \left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{A}^1 \rtimes \mathbb{G}_m^2}\right], \mu'\right)\right] = \frac{1}{2} \cdot \left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right], \mu_4' \circ i_1\right)\right] - \frac{3}{4} \cdot \left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{G}_m}\right], \mu_4' \circ i_2\right)\right] \\
& - \frac{1}{2} \cdot \left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{G}_m^2}\right], \mu' \circ i_1\right)\right] + \frac{1}{2} \left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{G}_m}\right], \mu' \circ i_2\right)\right]
\end{aligned} \tag{9.50}$$

which simplifies to:

$$\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}) = -\frac{1}{4} \left[\left(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{G}_m}\right], \mu' \circ i_2\right)\right]. \tag{9.51}$$

Now apply the Lie algebra homomorphism and obtain

$$\begin{aligned}&\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))=-\frac{1}{4}\cdot\chi^{na}\bigg(\left[\frac{\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}{\mathbb{G}_m}\right],(\mu'\circ i_2)^*\nu_{\mathfrak{M}_4^{(\beta,2)}}\bigg)\cdot\tilde{\lambda}^{(\beta,2)}\\&=(-1)^1\cdot(-1)^{\dim(\mathfrak{M}_4^{(\beta_k,\beta_l,3)}(\tilde{\tau}))-1}\cdot(-\frac{1}{4})\cdot\chi(\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau}))\cdot\tilde{\lambda}^{(\beta,2)}\end{aligned}\tag{9.52}$$

9.6 Computation of $\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{(\beta,2)}(\tilde{\tau}))$

In order to compute $\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{(\beta,2)}(\tilde{\tau}))$ in Equation (9.16) we add the contributions coming from $\epsilon_i^{(k,l)}$ (for all possible choices of β_k and β_l) to $\tilde{\Psi}^{\mathcal{B}_p}(\bar{\delta}_s^{(\beta,2)}(\tilde{\tau}))$, i.e:

$$\begin{aligned}&\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{(\beta,2)})(\tilde{\tau})=\sum_{\beta_k+\beta_l=\beta}^{o}\Big(\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_1^{(k,l)}(\tilde{\tau}))\Big)+\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_2^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))+\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau}))\\&+\sum_{\substack{\beta_k+\beta_l=\beta\\ \beta_k\neq\beta_l}}^{o}\Big(\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_3^{(k,l)}(\tilde{\tau}))\Big)+\tilde{\Psi}^{\mathcal{B}_p}(\epsilon_4^{(\frac{1}{2},\frac{1}{2})}(\tilde{\tau})).\end{aligned}\tag{9.53}$$

Let $d_i=\dim(\mathfrak{M}_i^{(\beta_k,\beta_l,3)}(\tilde{\tau}))-1$ for $i=1,\cdots,4$. By equations (9.23), (9.35),(9.45),(9.52) we obtain:

$$\begin{aligned}&\tilde{\Psi}^{\mathcal{B}_p}(\bar{\epsilon}^{(\beta,2)})(\tilde{\tau})=\chi^{na}(\left[\frac{\mathcal{M}_s^{(\beta,2)}(\tilde{\tau})}{\mathrm{GL}_2(\mathbb{C})}\right],\nu_{\mathfrak{M}_s^{(\beta,2)}})\cdot\tilde{\lambda}^{(\beta,2)}\\&+\sum_{\beta_k+\beta_l=\beta}^{o}\bigg(\frac{-1}{2}\cdot(-1)^{d_1}\Big(\int_{(E_1,E_3)\in\mathcal{M}^{(\beta_k,1)}(\tilde{\tau})\times\mathcal{M}^{(\beta_l,1)}(\tilde{\tau})\backslash\Delta}\chi(\mathbb{P}(\mathrm{Ext}^1(E_3,E_1)))d\chi\Big)\cdot\tilde{\lambda}^{(\beta,2)}\bigg)\\&-\frac{3}{2}\cdot(-1)^{d_2}\cdot\bigg(\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\chi\Big(\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))\Big)d\chi\bigg)\cdot\tilde{\lambda}^{(\beta,2)}\\&+\frac{1}{4}\cdot(-1)^{d_4}\cdot\chi(\mathcal{M}_s^{(\frac{\beta}{2},1)}(\tilde{\tau}))\cdot\tilde{\lambda}^{(\beta,2)}.\end{aligned}\tag{9.54}$$

10 Computation of the invariant

By the wall crossing computation of Joyce and Song in [9] (Equation 13.31):

$$\chi(\mathcal{M}_s^{(\frac{\beta}{2},1)}(\tilde{\tau})) = \sum_{1\leq l,\beta_1+\cdots+\beta_l=\frac{\beta}{2}} \Bigg[\frac{(1)}{l!}\cdot\bar{\chi}_{\mathcal{B}_p}((0,2),(\beta_1,0))\cdot$$
$$\prod_{i=1}^{l}\Bigg(\overline{DT}^{\beta_i}(\tau)\cdot\bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots+\beta_{i-1},2),(\beta_i,0))\cdot(-1)^{\bar{\chi}_{\mathcal{B}_p}((0,2),(\beta_1,0))+\sum_{i=1}^{l}\bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots\beta_{i-1},2),(\beta_i,0))}\Bigg)\Bigg]. \tag{10.1}$$

By Definition 5.7 and equations (10.1) and (9.54) we obtain:

$$\mathbf{B}_p^{ss}(X,\beta,2,\tilde{\tau}) =$$
$$\sum_{\beta_k+\beta_l=\beta}^{o}\Bigg[(-1)^{d_1+1}\cdot\frac{1}{2}\cdot\int_{(E_1,E_3)\in\mathfrak{G}^{(\beta_k,1)}\times\mathfrak{G}^{(\beta_l,1)}\backslash\Delta}\chi\Big(\mathbb{P}(\mathrm{Ext}^1(E_3,E_1))\Big)d\chi\Bigg]$$
$$+\frac{3}{2}\cdot(-1)^{d_2+1}\cdot\int_{E_1\in\mathcal{M}^{(\frac{\beta}{2},1)}(\tilde{\tau})}\chi\Big(\mathbb{P}(\mathrm{Ext}^1(E_1,E_1))\Big)d\chi$$
$$+\frac{1}{4}\cdot(-1)^{d_4}\cdot\sum_{1\leq l,\beta_1+\cdots+\beta_l=\frac{\beta}{2}}\Bigg[\frac{(1)}{l!}\cdot\bar{\chi}_{\mathcal{B}_p}((0,2),(\beta_1,0))\cdot\prod_{i=1}^{l}\Bigg(\overline{DT}^{\beta_i}(\tau)\cdot\bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots+\beta_{i-1},2),(\beta_i,0))$$
$$\cdot(-1)^{\bar{\chi}_{\mathcal{B}_p}((0,2),(\beta_1,0))+\sum_{i=1}^{l}\bar{\chi}_{\mathcal{B}_p}((\beta_1+\cdots\beta_{i-1},2),(\beta_i,0))}\Bigg)\Bigg]. \tag{10.2}$$

Remark 10.1. It is easy to see that by substituting $([\mathbb{P}^1],2)$ in Equation (10.2) one obtains the result obtained in (6.25) which is also compatible with the result of the wallcrossing computation obtained by substituting $([\mathbb{P}^1],2)$ in [15] (Equation 12.32).
One may also compute the right hand side of Equation (10.2) by substituting $(\beta,2)$ when β is not irreducible. For instance one can compute $\mathbf{B}_p^{ss}(X,2[\mathbb{P}^1],2,\tilde{\tau})$ where X is given by the total space of $\mathcal{O}_{\mathbb{P}^1}^{\oplus 2}(-1)\to\mathbb{P}^1$. This means to compute the invariant of $\tilde{\tau}$-semistable objects $(F,\mathbb{C}^2,\phi_{\mathbb{C}^2})$ of type $(2[\mathbb{P}^1],2)$ in $\mathcal{B}_p$. Assume $\chi(F)=r$. In this case, for a sheaf F appearing in such object of $\mathcal{B}_p$, if $r=2q+1$, then semistability implies stability and if $r=2q$ then F is given as a strictly semistable sheaf. Based on computations in [11] and [3] for $r=2q+1$ there exist no stable sheaves with $\mathrm{ch}_2(F)=2[\mathbb{P}^1]$. Now assume $r=2q$. In this case the semistable sheaves are given by $F=\mathcal{O}_{\mathbb{P}^1}(q-1)\oplus\mathcal{O}_{\mathbb{P}^1}(q-1)$. Note that a sheaf $F=\mathcal{O}_{\mathbb{P}^1}(a)\oplus\mathcal{O}_{\mathbb{P}^1}(b)$ where $a\neq b$ and $a+b=2q-2$ is not semistable since $a\neq q-1$ and $b\neq q-1$ and in that case either $\mathcal{O}_{\mathbb{P}^1}(b)$ or $\mathcal{O}_{\mathbb{P}^1}(a)$ destabilizes F. Now by substituting $(2[\mathbb{P}^1],2)$ in (10.2) and considering the sheaf F to satisfy the properties discussed above one obtains the invariant:

$$\mathbf{B}_p^{ss}(X,2[\mathbb{P}^1],2,\tilde{\tau}) = -\frac{1}{2}(n+q)^2-(n+q), \tag{10.3}$$

The computations in this case involve arguments similar to the ones given in Section 6 hence we have omitted the explicit calculations and therefore we leave them for the interested reader.

Remark 10.2. In [16] Toda has exploited similar stratification strategy for the moduli stack of objects composed of a zero dimensional sheaf F given as the quotient $\mathcal{O}_X^2 \twoheadrightarrow F$ where the objects are assumed to be semistable with respect to a stability condition in the sense of Bridgeland [1]. Moreover he has given an evidence of the integrality conjecture [12] (conjecture 6) for the corresponding partition functions associated to the moduli stack of these objects. The identities in [16] (equations 100, 99 and 88) play an important role in his proof of the integrality conjecture. Our stratification strategy and calculation of the invariants share many similarities with those in [16]. Equations (9.23) and (9.35) are analogue of equations 88 and 99 in [16] respectively and we suspect that in some cases they can be used to prove the integrality of the corresponding partition functions for the invariants of the moduli stack of objects in $\mathcal{B}_p$. Note that the stability condition used in our approach is the weak stability condition (Definition 3.23) used by Joyce-Song in [9] which, despite having a much simpler definition than the Bridgeland stability conditions, shares many strong properties with them. Finally, one can use Equation (10.2) and [15] (Corollary 13.7) to write the invariants of the higher rank Joyce-Song $\hat{\tau}$-limit stable pairs (Definition 3.11) in terms of the generalized Donaldson-Thomas invariants.

References

[1] T. Bridgeland. Stability conditions on triangulated categories. *Ann. of Math*, pages 317–345, 2007.

[2] T. Graber and R. Pandharipande. Localization of virtual classes. *Invent. Math.*, 135:487–518, 1999.

[3] Shinobu Hosono, Masa-Hiko Saito, and Atsushi Takahashi. Relative Lefschetz action and BPS state counting. *Int. Math. Res. Not.*, 2001, Issue 15:783–816, 2001.

[4] Dominic Joyce. Configurations in abelian categories. II. Ringel-Hall algebras. *arXiv:0503029v4*, 2005.

[5] Dominic Joyce. Motivic invariants of Artin stacks and stack functions. *Quarterly Journal of Mathematics*, 58 No 3:345–392, 2005.

[6] Dominic Joyce. Configurations in abelian categories. I. basic properties and moduli stacks. *Advances in Mathematics*, 203, Issue 1:194–255, 2006.

[7] Dominic Joyce. Configurations in abelian categories. III. Stability conditions and identities. *Advances in Mathematics*, 210, no. 2:635–706, 2007.

[8] Dominic Joyce. Configurations in abelian categories. IV. Invariants and changing stability conditions. *Advances in Mathematics*, 217, no. 1, 2008.

[9] Dominic Joyce and Yinan Song. A theory of generalized Donaldson-Thomas invariants. *arXiv:0810.5645*, 2009.

[10] Balazs Szendroi Kai Behrend, Jim Bryan. Motivic degree zero Donaldson-Thomas invariants. *arXiv:0909.5088v1*, 2009.

[11] Sheldon Katz. Genus zero Gopakumar-Vafa invariants of contractible curves. *J. Differential Geom.*, 79 Number 2:185–195, 2008.

[12] Yan Soibelman Maxim Kontsevich. Stability structures, motivic Donaldson-Thomas invariants and cluster transformations. *arXiv:0811.2435*, 2008.

[13] R. Pandharipande and R. P. Thomas. Curve counting via stable pairs in the derived category. *Inventiones*, 178:407–447, 2009.

[14] Artan Sheshmani. Higher rank stable pairs and virtual localization. *arXiv1011.6342*, 2010.

[15] Artan Sheshmani. Wall-crossing and invariants of higher rank stable pairs. *arXiv:1101.2252*, 2010.

[16] Yukinobu Toda. On a computation of rank two Donaldson-Thomas invariants. *arXiv:0912.2507*, 2010.